arXiv:1606.00183v2 [math.RA] 11 Jul 2016

THE CLASSIFICATION OF 3-DIMENSIONAL NOETHERIAN CUBIC CALABI-YAU ALGEBRAS

IZURU MORI AND KENTA UEYAMA

Abstract. It is known that every 3-dimensional noetherian Calabi-Yau algebra generated in degree 1 is isomorphic to a Jacobian algebra of a superpotential. Recently, S. P. Smith and the first author classified all superpotentials whose Jacobian algebras are 3-dimensional noetherian quadratic Calabi-Yau algebras. The main result of this paper is to classify all superpotentials whose Jacobian algebras are 3-dimensional noetherian cubic Calabi-Yau algebras. As an application, we show that if S is a 3-dimensional noetherian cubic Calabi-Yau algebra and σ is a graded algebra automorphism of S, then the homological determinant of σ can be calculated by the formula $\operatorname{hdet}\sigma = (\det\sigma)^2$ with one exception.

1. Introduction

Throughout this paper, we fix an algebraically closed field k of characteristic 0, and we assume that all vector spaces, algebras and unadorned tensor products are over k. In noncommutative algebraic geometry, AS-regular algebras are the most important class of algebras to study. In fact, the classification of 3-dimensional (noetherian) AS-regular algebras (generated in degree 1) using algebraic geometry is regarded as a starting point of noncommutative algebraic geometry ([1], [2]). On the other hand, in representation theory of algebras, Calabi-Yau algebras are important class of algebras to study. Since every connected graded Calabi-Yau algebra is AS-regular ([15]), it is interesting to study such algebras from the point of view of both noncommutative algebraic geometry and representation theory. In particular, since every m-Koszul Calabi-Yau algebra S is isomorphic to a Jacobian algebra $J(\mathsf{w}_S)$ of a unique superpotential w_S up to non-zero scalar multiples ([4], [5], [12]), it is interesting to study such algebras using both algebraic geometry and superpotentials. If S is a 3-dimensional noetherian Calabi-Yau algebra generated in degree 1 over k, then S is either 2-Koszul (quadratic) or 3-Koszul (cubic), so $S = J(\mathsf{w}_S)$ for some unique superpotential w_S. In [13], S. P. Smith and the first author of this paper focused on studying 3-dimensional noetherian quadratic Calabi-Yau algebras by these points of view. As a continuation, in this paper, we focus on studying 3-dimensional noetherian cubic Calabi-Yau algebras.

This paper is organized as follows: In section 2, we collect some preliminary results which are needed in this paper.

Section 3 is the heart of this paper. Let S be a 3-dimensional noetherian Calabi-Yau algebra. If S is quadratic, then $\mathsf{w}_S \in V^{\otimes 3}$ where V is a 3-dimensional vector space over k. In [13], all $\mathsf{w} \in V^{\otimes 3}$ such that $J(\mathsf{w})$ are 3-dimensional Calabi-Yau were classified. On the other hand, if S is cubic, then $\mathsf{w}_S \in V^{\otimes 4}$ where V is a 2-dimensional vector space over k. A natural next project is to classify all $\mathsf{w} \in V^{\otimes 4}$ such that $J(\mathsf{w})$ are 3-dimensional Calabi-Yau. The main result of this paper is to classify all such superpotentials. By this complete classification, we obtain the following results:

2010 *Mathematics Subject Classification.* 16E65, 16S38, 16W50.

Key words and phrases. Calabi-Yau algebras, superpotentials, homological determinants.

The first author was supported by JSPS Grant-in-Aid for Scientific Research (C) 25400037. The second author was supported by JSPS Grant-in-Aid for Young Scientists (B) 15K17503.

(1) We compute all possible point schemes for 3-dimensional noetherian cubic Calabi-Yau algebras (Theorem 3.6). By this computation, we see that not all bidegree (2, 2) divisors in $\mathbb{P}^1 \times \mathbb{P}^1$ appear as point schemes. This result contrasts to the fact that all degree 3 divisors in $\mathbb{P}^2$ appear as point schemes of 3-dimensional noetherian quadratic Calabi-Yau algebras ([13]).
(2) We show that $J(\mathsf{w})$ is 3-dimensional Calabi-Yau except for five algebras up to isomorphisms (Theorem 3.7).
(3) We show that $J(\mathsf{w})$ is 3-dimensional Calabi-Yau if and only if it is a domain (Corollary 3.8) as in the quadratic case ([13]).

Section 4 provides a further application of the classification. The homological determinant plays an important role in invariant theory for AS-regular algebras ([8], [10], [11], [6], [14] etc.). Contrary to its importance, it is rather mysterious and not easy to calculate from the definition. If $S = T(V)/(R)$ is a 3-dimensional noetherian quadratic Calabi-Yau algebra where $R \subset V \otimes V$, then it was shown in [12] that $\operatorname{hdet} \sigma = \det \sigma|_V$ for every $\sigma \in \operatorname{GrAut} S$ if and only if $\mathsf{w}_S \notin \operatorname{Sym}^3 V$. This means that for most of 3-dimensional noetherian quadratic Calabi-Yau algebras $S = T(V)/(R)$, $\operatorname{hdet} \sigma = \det \sigma|_V$ for every $\sigma \in \operatorname{GrAut} S$. By [9, Lemma 7.3 (2)], for typical examples of 3-dimensional noetherian cubic AS-regular algebras $S = T(V)/(R)$ where $R \subset V \otimes V \otimes V$, it holds that $\operatorname{hdet} \sigma = (\det \sigma|_V)^2$ for every $\sigma \in \operatorname{GrAut} S$, so it is natural to expect that, for a 3-dimensional noetherian cubic Calabi-Yau algebra $S = T(V)/(R)$, $\operatorname{hdet} \sigma = (\det \sigma|_V)^2$ for every $\sigma \in \operatorname{GrAut} S$ if and only if $\mathsf{w}_S \notin \operatorname{Sym}^4 V$. Although one direction of the above expectation is true (Corollary 4.6), we will show a slightly more striking result, namely, for a 3-dimensional noetherian cubic Calabi-Yau algebra $S = T(V)/(R)$, $\operatorname{hdet} \sigma = (\det \sigma|_V)^2$ for every $\sigma \in \operatorname{GrAut} S$ if and only if

$$S \ncong k\langle x, y\rangle/(xy^2 + yxy + y^2x + \sqrt{-3}x^3, yx^2 + xyx + x^2y + \sqrt{-3}y^3)$$

(Theorem 4.8).

For a 3-dimensional noetherian quadratic Calabi-Yau algebra $S = T(V)/(R)$, it was also shown in [13] that $\mathsf{w}_S \notin \operatorname{Sym}^3 V$ if and only if S is a deformation quantization of $k[x, y, z]$, and $\mathsf{w}_S \in \operatorname{Sym}^3 V$ if and only if S is a Clifford algebra. In the last section, we will show that something similar holds in one direction (Theorem 5.1, Theorem 5.3).

2. Preliminaries

Throughout this paper, let k be an algebraically closed field of characteristic 0, and V a finite dimensional vector space over k. In this section, we collect some preliminary results which are needed in this paper.

2.1. **Superpotentials.**

We define the action of $\theta \in \mathfrak{S}_m$ on $V^{\otimes m}$ by

$$\theta(v_1 \otimes \cdots \otimes v_m) := v_{\theta(1)} \otimes \cdots \otimes v_{\theta(m)}.$$

Specializing to the m-cycle $\phi \in \mathfrak{S}_m$, we define

$$\phi(v_1 \otimes v_2 \otimes \cdots \otimes v_{m-1} \otimes v_m) := v_m \otimes v_1 \otimes \cdots \otimes v_{m-2} \otimes v_{m-1}.$$

We define linear maps $c, s, a : V^{\otimes m} \to V^{\otimes m}$ by

$$c(\mathsf{w}) := \frac{1}{m}\sum_{i=0}^{m-1} \phi^i(\mathsf{w}),$$
$$s(\mathsf{w}) := \frac{1}{|\mathfrak{S}_m|}\sum_{\theta\in\mathfrak{S}_m} \theta(\mathsf{w}),$$
$$a(\mathsf{w}) := \frac{1}{|\mathfrak{S}_m|}\sum_{\theta\in\mathfrak{S}_m} (\operatorname{sgn}\theta)\theta(\mathsf{w}).$$

We define the following subspaces of $V^{\otimes m}$:

$$\operatorname{Sym}^m V := \{\mathsf{w} \in V^{\otimes m} \mid \theta(\mathsf{w}) = \mathsf{w} \text{ for all } \theta \in \mathfrak{S}_m\},$$
$$\operatorname{Alt}^m V := \{\mathsf{w} \in V^{\otimes m} \mid \theta(\mathsf{w}) = (\operatorname{sgn}\theta)\mathsf{w} \text{ for all } \theta \in \mathfrak{S}_m\}.$$

It is easy to see that $\operatorname{Sym}^m V = \operatorname{Im} s$ and $\operatorname{Alt}^m V = \operatorname{Im} a$. In this paper, $\operatorname{Im} c$ also plays an important role.

Definition 2.1. Let $\mathsf{w} \in V^{\otimes m}$ and $\sigma \in \mathrm{GL}(V)$. We call w

(1) a superpotential if $\phi(\mathsf{w}) = \mathsf{w}$,
(2) a σ-twisted superpotential if $(\sigma \otimes \mathrm{id}^{\otimes m-1})\phi(\mathsf{w}) = \mathsf{w}$,
(3) a twisted superpotential if it is a σ-twisted superpotential for some σ.

Lemma 2.2. [12, Lemma 6.1] *For $\mathsf{w} \in V^{\otimes m}$, w is a superpotential if and only if $\mathsf{w} \in \operatorname{Im} c$.*

We call $\operatorname{Im} c$ the space of superpotentials. For $\sigma \in \mathrm{GL}(V)$ and $\mathsf{w} \in V^{\otimes m}$, we often write $\sigma(\mathsf{w}) := \sigma^{\otimes m}(\mathsf{w})$ by abuse of notation.

Lemma 2.3. *For $\theta \in \mathfrak{S}_m, \sigma \in \mathrm{GL}(V), \mathsf{w} \in V^{\otimes m}$, we have*

(1) $\theta(\sigma(\mathsf{w})) = \sigma(\theta(\mathsf{w}))$,
(2) $c(\sigma(\mathsf{w})) = \sigma(c(\mathsf{w}))$,
(3) $s(\sigma(\mathsf{w})) = \sigma(s(\mathsf{w}))$,
(4) $a(\sigma(\mathsf{w})) = \sigma(a(\mathsf{w}))$.

Proof. For $v_1 \otimes \cdots \otimes v_m \in V^{\otimes m}$,

$$\begin{aligned}\theta(\sigma(v_1 \otimes \cdots \otimes v_m)) &= \theta(\sigma(v_1) \otimes \cdots \otimes \sigma(v_m)) = \sigma(v_{\theta(1)}) \otimes \cdots \otimes \sigma(v_{\theta(m)}) \\ &= \sigma(v_{\theta(1)} \otimes \cdots \otimes v_{\theta(m)}) = \sigma(\theta(v_1 \otimes \cdots \otimes v_m)),\end{aligned}$$

so $\theta(\sigma(\mathsf{w})) = \sigma(\theta(\mathsf{w}))$. It follows that

$$c(\sigma(\mathsf{w})) = \frac{1}{m}\sum_{i=0}^{m-1} \phi^i(\sigma(\mathsf{w})) = \frac{1}{m}\sum_{i=0}^{m-1} \sigma(\phi^i(\mathsf{w})) = \sigma\left(\frac{1}{m}\sum_{i=0}^{m-1} \phi^i(\mathsf{w})\right) = \sigma(c(\mathsf{w})).$$

Similarly, we have $s(\sigma(\mathsf{w})) = \sigma(s(\mathsf{w}))$ and $a(\sigma(\mathsf{w})) = \sigma(a(\mathsf{w}))$. □

Let V be a finite dimensional vector space over k. The tensor algebra of V over k is denoted by $T(V)$, which is an $\mathbb{N}$-graded algebra by $T(V)_i = V^{\otimes i}$. An m-homogeneous algebra is an $\mathbb{N}$-graded algebra of the form $T(V)/(R)$ where (R) is the two-sided ideal generated by a subspace $R \subset V^{\otimes m}$. The symmetric algebra of V over k is denoted by $S(V) = T(V)/(R)$ where $R = \{u \otimes v - v \otimes u \in V \otimes V \mid u, v \in V\}$, which is an example of a 2-homogeneous algebra (a quadratic algebra). Note that $S(V)_m = V^{\otimes m}/\sum_{i+j+2=m} V^i \otimes R \otimes V^j$ is the quotient space. We denote the quotient map by $\overline{(-)} : V^{\otimes m} \to S(V)_m$. Since $s(\mathsf{w}) = 0$ for every $\mathsf{w} \in V^i \otimes R \otimes V^j$, the linear map $s : V^{\otimes m} \to V^{\otimes m}$ induces a linear map $\widetilde{(-)} : S(V)_m \to V^{\otimes m}$, called the symmetrization map.

Lemma 2.4. *The linear maps* $\overline{(-)} : V^{\otimes m} \to S(V)_m$ *and* $\widetilde{(-)} : S(V)_m \to V^{\otimes m}$ *induce isomorphisms* $\overline{(-)} : \operatorname{Sym}^m V \to S(V)_m$ *and* $\widetilde{(-)} : S(V)_m \to \operatorname{Sym}^m V$ *which are inverses to each other.*

Lemma 2.5. *Let* $\sigma \in \mathrm{GL}(V)$.

(1) *For* $\mathsf{w} \in V^{\otimes m}$, $\sigma(\overline{\mathsf{w}}) = \overline{\sigma(\mathsf{w})}$.
(2) *For* $f \in S(V)_m$, $\sigma\left(\widetilde{f}\right) = \widetilde{\sigma(f)}$.

Proof. (1) This follows from the fact that σ extends to a graded algebra automorphism $\sigma \in \operatorname{GrAut} T(V)$, which induces a graded algebra automorphism $\sigma \in \operatorname{GrAut} S(V)$ by the formula $\sigma(\overline{\mathsf{w}}) = \overline{\sigma(\mathsf{w})}$.

(2) For every $f \in S(V)_m$, there exists $\mathsf{w} \in V^{\otimes m}$ such that $f = \overline{\mathsf{w}}$, so

$$\sigma\left(\widetilde{f}\right) = \sigma\left(\widetilde{\overline{\mathsf{w}}}\right) = \sigma(s(\mathsf{w})) = s(\sigma(\mathsf{w})) = \widetilde{\overline{\sigma(\mathsf{w})}} = \widetilde{\sigma(\overline{\mathsf{w}})} = \widetilde{\sigma(f)}$$

by Lemma 2.3 and (1). □

2.2. **Calabi-Yau Algebras.** Let V be a vector space, $W \subset V^{\otimes m}$ a subspace and $\mathsf{w} \in V^{\otimes m}$. We introduce the following notation:

$$\begin{aligned}
\partial W &:= \{(\psi \otimes \mathrm{id}^{\otimes m-1})(\mathsf{w}) \mid \psi \in V^*, \mathsf{w} \in W\},\\
W\partial &:= \{(\mathrm{id}^{\otimes m-1} \otimes \psi)(\mathsf{w}) \mid \psi \in V^*, \mathsf{w} \in W\},\\
\mathcal{D}(W, i) &:= T(V)/(\partial^i W),\\
\mathcal{D}(\mathsf{w}) &:= \mathcal{D}(k\mathsf{w}, 1),\\
J(\mathsf{w}) &:= \mathcal{D}(c(\mathsf{w})).
\end{aligned}$$

We call $J(\mathsf{w})$ the Jacobian algebra of w, and w the potential of $J(\mathsf{w})$. Note that $\mathcal{D}(\mathsf{w})$ and $J(\mathsf{w})$ are $(m-1)$ homogeneous algebras.

Choose a basis $x_1, \dots, x_n$ for V so that $T(V) = k\langle x_1, \dots, x_n\rangle$ and $S(V) = k[x_1, \dots, x_n]$. For $f \in k[x_1, \dots, x_n]$, the usual partial derivative with respect to x_i is denoted by f_{x_i}. For a monomial $\mathsf{w} = x_{i_1}x_{i_2}\cdots x_{i_{m-1}}x_{i_m} \in k\langle x_1, \dots, x_n\rangle_m$ of degree m, we define

$$\partial_{x_i}\mathsf{w} := \begin{cases} x_{i_2}\cdots x_{i_{m-1}}x_{i_m} & \text{if } i_1 = i \\ 0 & \text{if } i_1 \neq i, \end{cases} \quad \text{and} \quad \mathsf{w}\partial_{x_i} := \begin{cases} x_{i_1}x_{i_2}\cdots x_{i_{m-1}} & \text{if } i_m = i \\ 0 & \text{if } i_m \neq i. \end{cases}$$

We extend the maps $\partial_{x_i} : k\langle x_1, \dots, x_n\rangle_m \to k\langle x_1, \dots, x_n\rangle_{m-1}$ by linearity. In this notation, $\mathcal{D}(\mathsf{w}) = k\langle x_1, \dots, x_n\rangle/(\partial_{x_1}\mathsf{w}, \dots, \partial_{x_n}\mathsf{w})$.

Lemma 2.6. *Let* V *be a vector space spanned by* $x_1, \dots, x_n$.

(1) *For every* $f \in S(V)_m$, $\widetilde{f_{x_i}} = m\partial_{x_i}\widetilde{f}$.
(2) *For every* $\mathsf{w} \in \operatorname{Sym}^m V$, $\widetilde{\overline{\mathsf{w}}_{x_i}} = m\partial_{x_i}\mathsf{w}$.

Proof. This follows from [13, Lemma 2.5]. □

For an $\mathbb{N}$-graded algebra A, we denote by A^o the opposite graded algebra of A, and $A^e = A^o \otimes A$ the enveloping algebra of A. For graded left A-modules M, N, we denote by $\operatorname{Ext}^i_A(M, N)$ the i-th derived functor of $\operatorname{Hom}_A(M, N)$ in the category of graded left A-modules, and $\underline{\operatorname{Ext}}^i_A(M, N) := \bigoplus_{j\in\mathbb{Z}} \operatorname{Ext}^i_A(M, N(j))$ where $N(j) = N$ as a left A-module with the new grading $N(j)_n = N_{j+n}$. A connected graded algebra is an $\mathbb{N}$-graded algebra such that $A_0 = k$. In this case, we view $k = A/A_{\geq 1}$ as a graded A-module.

Definition 2.7. A connected graded algebra S is called a d-dimensional AS-regular algebra if

(1) $\operatorname{gldim} S = d < \infty$, and

(2) for some $\ell \in \mathbb{Z}$, $\underline{\mathrm{Ext}}^i_S(k,S) \cong \begin{cases} k(\ell) & \text{if } i = d, \\ 0 & \text{otherwise.} \end{cases}$

The integer ℓ is called the Gorenstein parameter of S.

Definition 2.8. An $\mathbb{N}$-graded algebra S is called a d-dimensional Calabi-Yau if

(1) S has a resolution of finite length by finitely generated graded projective left S^e-modules, and

(2) for some $\ell \in \mathbb{Z}$, $\underline{\mathrm{Ext}}^i_{S^e}(S,S^e) \cong \begin{cases} S(\ell) & \text{if } i = d, \\ 0 & \text{otherwise} \end{cases}$ as graded right A^e-modules.

It is known that a connected graded d-dimensional Calabi-Yau algebra is a d-dimensional AS-regular algebra ([15, Lemma 1.2]).

Theorem 2.9. [12, Proposition 2.12] *If S is a 3-dimensional noetherian AS-regular algebra generated in degree 1 over k, then there exists a unique twisted superpotential w_S up to non-zero scalar multiples such that $S = \mathcal{D}(\mathsf{w}_S)$. Moreover, if S is Calabi-Yau, then there exists a unique superpotential w_S up to non-zero scalar multiples such that $S = J(\mathsf{w}_S)$.*

Lemma 2.10. *Let $\sigma \in \mathrm{GL}(V)$ and $W \subset V^{\otimes m}$ a subspace. If $W' = \sigma^{\otimes m}(W) \subset V^{\otimes m}$, then σ extends to an isomorphism of graded algebras $\mathcal{D}(W,i) \to \mathcal{D}(W',i)$ for every $i \in \mathbb{N}$.*

Proof. Note that

$$\begin{aligned} \sigma^{\otimes m-1}(\partial W) &= \{\sigma^{\otimes m-1}(\psi \otimes \mathrm{id}^{\otimes m-1})(\mathsf{w}) \mid \psi \in V^*, \mathsf{w} \in W\} = \{(\psi \otimes \sigma^{\otimes m-1})(\mathsf{w}) \mid \psi \in V^*, \mathsf{w} \in W\} \\ &= \{(\psi\sigma \otimes \sigma^{\otimes m-1})(\mathsf{w}) \mid \psi \in V^*, \mathsf{w} \in W\} = \{(\psi \otimes \mathrm{id}^{\otimes m-1})\sigma^{\otimes m}(\mathsf{w}) \mid \psi \in V^*, \mathsf{w} \in W\} \\ &= \partial(\sigma^{\otimes m}(W)). \end{aligned}$$

Since $\sigma^{\otimes m}(W) = W'$, $\sigma^{\otimes m-i}(\partial^i W) = \partial^i(\sigma^{\otimes m}(W)) = \partial^i W'$ for every $i \in \mathbb{N}$ by induction, so σ extends to an isomorphism of graded algebras $\mathcal{D}(W,i) = T(V)/(\partial^i W) \to \mathcal{D}(W',i) = T(V)/(\partial^i W')$. □

A (super)potential $\mathsf{w} \in V^{\otimes m}$ is called Calabi-Yau if $J(\mathsf{w})$ is Calabi-Yau. Two potentials $\mathsf{w}, \mathsf{w}' \in V^{\otimes m}$ are called equivalent, denoted by $\mathsf{w} \sim \mathsf{w}'$, if $\mathsf{w}' = \sigma(\mathsf{w})$ for some $\sigma \in \mathrm{GL}(V)$.

Theorem 2.11. *Let $S = T(V)/(R), S' = T(V)/(R')$ be m-Koszul d-dimensional AS-regular algebras with the Gorenstein parameter ℓ. Then $S \cong S'$ as graded algebras if and only if $\mathsf{w}_S \sim \mathsf{w}_{S'}$. In particular, let $\mathsf{w}, \mathsf{w}' \in \mathrm{Im}\, c \subset V^{\otimes m}$ be Calabi-Yau superpotentials. Then $J(\mathsf{w}) \cong J(\mathsf{w}')$ as graded algebras if and only if $\mathsf{w} \sim \mathsf{w}'$.*

Proof. If $\phi : S = T(V)/(R) \to S' = T(V)/(R')$ is an isomorphism of graded algebras, then $\sigma := \phi|_V \in \mathrm{GL}(V)$ has the property that $\sigma(R) = R'$. By [12, Proposition 2.12],

$$\sigma(k\mathsf{w}_S) = \sigma\left(\bigcap_{i+j+m=\ell} V^{\otimes s} \otimes R \otimes V^{\otimes t}\right) = \bigcap_{i+j+m=\ell} V^{\otimes s} \otimes R' \otimes V^{\otimes t} = k\mathsf{w}_{S'},$$

so $\mathsf{w}_{S'} = \sigma(\mathsf{w}_S)$ by adjusting the scalar.

Conversely, if $\mathsf{w}_{S'} = \sigma(\mathsf{w}_S)$ for some $\sigma \in \mathrm{GL}(V)$, then $\sigma(k\mathsf{w}_S) = k\mathsf{w}_{S'}$, so σ extends to an isomorphism of graded algebras $S = \mathcal{D}(k\mathsf{w}_S, \ell - m) \to S' = \mathcal{D}(k\mathsf{w}_{S'}, \ell - m)$ by [12, Theorem 1.9] and Lemma 2.10. □

In [2], Artin, Tate and Van den Bergh classified of 3-dimensional noetherian AS-regular algebras generated in degree 1 using algebraic geometry. They showed that every 3-dimensional noetherian quadratic AS-regular algebra determines, and is determined by, a geometric triple $(E, \tau, \mathcal{L})$ where E is $\mathbb{P}^2$ or a divisor in $\mathbb{P}^2$ of degree 3, τ is an automorphism of E, and $\mathcal{L}$ is an invertible sheaf on

E, and also showed that every 3-dimensional noetherian cubic AS-regular algebra determines, and is determined by, a geometric triple $(E, \tau, \mathcal{L})$ where E is $\mathbb{P}^1 \times \mathbb{P}^1$ or a divisor in $\mathbb{P}^1 \times \mathbb{P}^1$ of bidegree (2, 2), τ is an automorphism of E, and $\mathcal{L}$ is an invertible sheaf on E. The scheme E is called the point scheme of S, and the pair (E, τ) is called the geometric pair of S. See [2] for details of geometric triples.

3. Classification of Calabi-Yau Superpotentials

In this section, we classify all superpotentials w such that $J(\mathsf{w})$ is 3-dimensional noetherian cubic Calabi-Yau up to isomorphisms of $J(\mathsf{w})$.

3.1. **Calabi-Yau Properties.** Let $S = T(V)/(R)$ be an m-homogeneous algebra such that $\dim V = \dim R = n$. If we choose a basis $x_1, \dots, x_n$ for V and a basis $f_1, \dots, f_n$ for R, then we may write $f_i = \sum_{j=1}^n m_{ij}x_j$ for some $n \times n$ matrix $\mathbf{M} = (m_{ij})$ whose entries are in $V^{\otimes m-1}$. Define $g_j = \sum_{i=1}^n x_i m_{ij}$. If we write $\mathbf{x} = \begin{pmatrix} x_1 \\ \vdots \\ x_n \end{pmatrix}, \mathbf{f} = \begin{pmatrix} f_1 \\ \vdots \\ f_n \end{pmatrix}, \mathbf{g} = \begin{pmatrix} g_1 \\ \vdots \\ g_n \end{pmatrix}$, then $\mathbf{f} = \mathbf{M}\mathbf{x}$, $\mathbf{g} = (\mathbf{x}^T\mathbf{M})^T$.

Remark 3.1. Note that $(\mathbf{x}^T\mathbf{M})^T \neq \mathbf{M}^T\mathbf{x}$.

Definition 3.2. [2] An m-homogeneous algebra $S = T(V)/(R)$ is called standard if $\dim V = \dim R = n$ and there exist a choice of a basis $x_1, \dots, x_n$ for V and a choice of a basis $f_1, \dots, f_n$ for R such that $\mathbf{g} = Q_S\mathbf{f}$ for some $Q_S \in \mathrm{GL}(n, k)$.

Fix a basis $x_1, \dots, x_n$ for V. For $\mathsf{w} \in V^{\otimes m}$, we define the $n \times n$ matrix $\mathbf{M}(\mathsf{w}) := (\partial_{x_i}\mathsf{w}\partial_{x_j})$ whose entries are in $V^{\otimes m-2}$. If we write $\mathbf{x} = \begin{pmatrix} x_1 \\ \vdots \\ x_n \end{pmatrix}$, then $\mathbf{M}(\mathsf{w})$ is the unique matrix such that $\mathsf{w} = \mathbf{x}^T\mathbf{M}(\mathsf{w})\mathbf{x}$. Viewing $\mathbf{M}(\mathsf{w}) \in M_n(S(V)^{\circ m-2})$, we define the (noncommutative) Hessian of w by $H(\mathsf{w}) = \det \mathbf{M}(\mathsf{w}) = \det(\partial_{x_i}\mathsf{w}\partial_{x_j}) \in S(V)^{\circ m-2}$ where $A \circ B$ denotes the Segre product of A and B.

Let V be a vector space spanned by $x_1, \dots, x_n$ and $\mathsf{w} \in V^{\otimes m}$. Since $J(\mathsf{w}) := T(V)/(\partial_{x_1}c(\mathsf{w}), \dots, \partial_{x_n}c(\mathsf{w}))$, and $(\partial_{x_1}c(\mathsf{w}) \ \cdots \ \partial_{x_n}c(\mathsf{w}))^T = \mathbf{M}(c(\mathsf{w}))\mathbf{x}$, we have the following results:

Proposition 3.3. *Let V be a vector space spanned by $x_1, \dots, x_n$ and $\mathsf{w} \in V^{\otimes m}$. Then $J(\mathsf{w})$ is standard if and only if $\partial_{x_1}c(\mathsf{w}), \dots, \partial_{x_n}c(\mathsf{w})$ are linearly independent.*

Proof. This follows from [13, Proposition 2.6]. □

Theorem 3.4. *Let V be a 2-dimensional vector space spanned by x, y and $\mathsf{w} \in V^{\otimes 4}$. Then $J(\mathsf{w})$ is 3-dimensional Calabi-Yau if and only if $\partial_x c(\mathsf{w}), \partial_y c(\mathsf{w})$ are linearly independent and*

$$\mathcal{V}(\partial_x c(\mathsf{w})\partial_x, \partial_x c(\mathsf{w})\partial_y, \partial_y c(\mathsf{w})\partial_x, \partial_y c(\mathsf{w})\partial_y) = \emptyset$$

in $\mathbb{P}^1 \times \mathbb{P}^1$. If this is the case, then the point scheme of $J(\mathsf{w})$ is given by $\mathcal{V}(H(c(\mathsf{w})))$.

Proof. This follows from Proposition 3.3 and [2, Theorem 1]. □

3.2. **Classification of four points in $\mathbb{P}^1$.** Two polynomials $f, f' \in S(V)_m$ are called equivalent, denoted by $f \sim f'$, if $f' = \sigma(f)$ for some $\sigma \in \mathrm{GL}(V)$. Note that $f \sim f'$ if and only if $\operatorname{Proj} S(V)/(f)$ and $\operatorname{Proj} S(V)/(f')$ are projectively equivalent, denoted by $\operatorname{Proj} S(V)/(f) \sim \operatorname{Proj} S(V)/(f')$.

Lemma 3.5. *Let V be a 2-dimensional vector space spanned by x, y. Every $f \in S(V)_4$ is equivalent to one of the following:*

(1) $f_1 = 0$;
(2) $f_2 = x^4$;
(3) $f_3 = x^3y$;

(4) $f_4 = x^2y^2$;
(5) $f_5 = x^4 + x^2y^2$;
(6) $g_\lambda = x^4 + y^4 + \lambda x^2y^2, \lambda \in k \setminus \{\pm 2\}$.

Moreover, in case (6), $g_\lambda \sim g_{\lambda'}$ if and only if $\lambda' = \pm\lambda$ or $(2 \pm \lambda)(2 \pm \lambda') = 16$ or $(2 \pm \lambda)(2 \mp \lambda') = 16$.

Proof. It is equivalent to classify $X := \operatorname{Proj} k[x, y]/(f)$ up to projective equivalences. Note that $f \neq 0$ if and only if $X \neq \mathbb{P}^1$, and, in this case, $X = \operatorname{Proj} k[x, y]/(f)$ is a set of at most four points. Note also that every sets of three points (with fixed multiplicities) are projectively equivalent to each other. If $X = \mathbb{P}^1$, then $X = \operatorname{Proj} k[x, y]$. If X consists of one point $(0, 1)$ with multiplicity 4, then $X = \operatorname{Proj} k[x, y]/(x^4)$. If X consists of a point $(0, 1)$ with multiplicity 3 and a point $(1, 0)$ of multiplicity 1, then $X = \operatorname{Proj} k[x, y]/(x^3y)$. If X consists of two points $(0, 1), (1, 0)$ with multiplicity 2, then $X = \operatorname{Proj} k[x, y]/(x^2y^2)$. If X consists of a point $(0, 1)$ with multiplicity 2 and two points $(1, \sqrt{-1}), (1, -\sqrt{-1})$ of multiplicity 1, then $X = \operatorname{Proj} k[x, y]/(x^2(x^2 + y^2))$. If $X = \operatorname{Proj} k[x, y]/(f)$ consists of four points $(0, 1), (1, 0), (1, -1), (1, -\lambda)$ of multiplicity 1 where $\lambda \neq 0, 1$, then

$$f = xy(x + y)(\lambda x + y) \sim x^3y + (\frac{1}{\sqrt{\lambda}} + \sqrt{\lambda})x^2y^2 + xy^3.$$

Since $\lambda \neq 0, 1$, we have $\frac{1}{\sqrt{\lambda}} + \sqrt{\lambda} \neq \pm 2$, so we may write $f \sim xy(x^2 + y^2 + \lambda xy)$ where $\lambda \neq \pm 2$. Moreover, one can check that

$$f \sim xy(x^2 + y^2 + \lambda xy) \sim x^4 + y^4 - \frac{2\lambda}{\sqrt{\lambda^2 - 4}}x^2y^2.$$

It is easy to see that $-\frac{2\lambda}{\sqrt{\lambda^2-4}} \neq \pm 2$, so we may write $f \sim x^4 + y^4 + \lambda x^2y^2$ where $\lambda \neq \pm 2$. On the other hand, if $\lambda \neq \pm 2$, then it is easy to see that $\operatorname{Proj} k[x, y]/(x^4 + y^4 + \lambda x^2y^2)$ consists of four distinct points.

Write $g_\lambda := x^4 + y^4 + \lambda x^2y^2$. Let $\sigma \in \mathrm{GL}(2, k)$ such that $\sigma(g_\lambda) = g_{\lambda'}$ for some $\lambda, \lambda' \in k \setminus \{\pm 2\}$. By a direct calculation, we see that σ is given by

$$\begin{pmatrix} \alpha & 0 \\ 0 & \delta \end{pmatrix} (\delta^4 = \alpha^4), \quad \begin{pmatrix} 0 & \beta \\ \gamma & 0 \end{pmatrix} (\gamma^4 = \beta^4), \text{ or } \begin{pmatrix} \alpha & \beta \\ -\xi\alpha & \xi\beta \end{pmatrix} (\beta^4 = \alpha^4, \xi : \text{4-th root of unity}).$$

If σ is in the first case or the second case, then we have $\lambda' = \pm\lambda$. If $\sigma = \left(\begin{smallmatrix} \alpha & \beta \\ -\xi\alpha & \xi\beta \end{smallmatrix}\right)$ with $\beta^2 = \alpha^2, \xi^2 = 1$, then

$$(\alpha x + \beta y)^4 + (-\xi\alpha x + \xi\beta y)^4 + \lambda(\alpha x + \beta y)^2(-\xi\alpha x + \xi\beta y)^2 = \alpha^4(2 + \lambda)\left(x^4 + y^4 + \frac{12 - 2\lambda}{2 + \lambda}x^2y^2\right),$$

so we have $\lambda' = \frac{12-2\lambda}{2+\lambda}$. Thus $(2 + \lambda')(2 + \lambda) = 16$. By similar calculations, we can show that if $\beta^2 = -\alpha^2, \xi^2 = 1$, then $(2 - \lambda')(2 + \lambda) = 16$, and if $\beta^2 = \pm\alpha^2, \xi^2 = -1$, then $(2 \pm \lambda')(2 - \lambda) = 16$, so the assertion holds. □

3.3. **Classification of Calabi-Yau Superpotentials.** Let V be a 2-dimensional vector space spanned by x, y. It is clear that the space of superpotentials $\operatorname{Im} c \subset V^{\otimes 4}$ has a basis

$$\begin{aligned} \mathsf{w}_1 &= x^2y^2 + xy^2x + y^2x^2 + yx^2y, \\ \mathsf{w}_2 &= xyxy + yxyx, \\ \mathsf{w}_3 &= x^3y + x^2yx + xyx^2 + yx^3, \\ \mathsf{w}_4 &= y^3x + y^2xy + yxy^2 + xy^3, \\ \mathsf{w}_5 &= x^4, \\ \mathsf{w}_6 &= y^4. \end{aligned}$$

For the rest of this paper, we fix the above basis for $\operatorname{Im} c$. It is easy to see that $\operatorname{Sym}^4 V$ is a codimension 1 subspace of $\operatorname{Im} c$ spanned by $\mathsf{w}_1+\mathsf{w}_2, \mathsf{w}_3, \mathsf{w}_4, \mathsf{w}_5, \mathsf{w}_6$. We set $W' := \bigoplus_{i=3}^{6} k\mathsf{w}_i \subset \operatorname{Sym}^4 V$, so that every $\mathsf{w} \in \operatorname{Im} c$ can be uniquely written as $\mathsf{w} = \alpha\mathsf{w}_1 + \beta\mathsf{w}_2 + \mathsf{w}'$ where $\alpha, \beta \in k$ and $\mathsf{w}' \in W'$.

Let $\mathsf{w}, \mathsf{w}' \in \operatorname{Im} c$ be Calabi-Yau superpotentials. By Theorem 2.11 and Lemma 2.5, if $J(\mathsf{w}) \cong J(\mathsf{w}')$ as graded algebras, then $\overline{\mathsf{w}} \sim \overline{\mathsf{w}'}$, so we may assume that $\overline{\mathsf{w}} \in S(V)_4$ is one of the forms in Lemma 3.5 to classify Calabi-Yau superpotentials $\mathsf{w} \in \operatorname{Im} c$ up to isomorphisms of $J(\mathsf{w})$.

Theorem 3.6. *Let V be a 2-dimensional vector space. The table below (Table 1) gives a classification of all Calabi-Yau superpotentials $0 \neq \mathsf{w} \in V^{\otimes 4}$ up to isomorphisms of $J(\mathsf{w})$. In each Calabi-Yau superpotential w, the defining relations of $J(\mathsf{w})$, and the geometric pair of $J(\mathsf{w})$ are also listed in the table. (In the table, a curve of bidegree (a,b) in $\mathbb{P}^1\times\mathbb{P}^1$ is denoted by (a,b) by abuse of notations.)*

Proof. Let V be a 2-dimensional vector space spanned by x, y and $\mathsf{w} \in \operatorname{Im} c$ a superpotential. We divide the cases according to the classification of $\overline{\mathsf{w}} \in S(V)_4$ as in Lemma 3.5.

(Case 1) If $\overline{\mathsf{w}} = f_1 = 0$, then $\mathsf{w} = \alpha\mathsf{w}_1 + \beta\mathsf{w}_2$ for $\alpha, \beta \in k$ such that $4\alpha + 2\beta = 0$. Since $\mathsf{w} \neq 0$, we have $\alpha \neq 0$, so we may assume that $\alpha = 1$. Since

$$\partial_x \mathsf{w} = xy^2 + y^2x - 2yxy, \quad \partial_y \mathsf{w} = yx^2 + x^2y - 2xyx$$

are linearly independent, $J(\mathsf{w})$ is standard by Proposition 3.3. It is easy to see that $\mathbf{M}(\mathsf{w}) = \begin{pmatrix} y^2 & xy-2yx \\ yx-2xy & x^2 \end{pmatrix}$. Moreover we can calculate $\mathcal{V}(y^2, xy-2yx, yx-2xy, x^2) = \emptyset$ in $\mathbb{P}^1\times\mathbb{P}^1$. Thus $J(\mathsf{w})$ is 3-dimensional Calabi-Yau by Theorem 3.4. The point scheme E of $J(\mathsf{w})$ is given by

$$\begin{aligned}\mathcal{V}(H(\mathsf{w})) &= \mathcal{V}\left(\det\begin{pmatrix} y_1y_2 & x_1y_2-2y_1x_2 \\ y_1x_2-2x_1y_2 & x_1x_2 \end{pmatrix}\right) \\ &= \mathcal{V}((x_1y_2-y_1x_2)^2) \subset \mathbb{P}^1\times\mathbb{P}^1\end{aligned}$$

(a double curve of bidegree (1, 1)), and the automorphism $\tau \in \operatorname{Aut}_k E_{\mathrm{red}}$ is given by

$$\tau((p_1,q_1),(p_1,q_1)) = ((p_1,q_1),(p_1,q_1)).$$

(Case 2) If $\overline{\mathsf{w}} = f_2 = x^4$, then $\mathsf{w} = \alpha\mathsf{w}_1 + \beta\mathsf{w}_2 + \mathsf{w}_5$ for $\alpha, \beta \in k$ such that $4\alpha + 2\beta = 0$. We have

$$\partial_x \mathsf{w} = \alpha(xy^2 + y^2x - 2yxy) + x^3, \quad \partial_y \mathsf{w} = \alpha(yx^2 + x^2y - 2xyx).$$

If $\alpha = 0$, then

$$\partial_x \mathsf{w} = x^3, \quad \partial_y \mathsf{w} = 0,$$

so $J(\mathsf{w})$ is not 3-dimensional Calabi-Yau by Theorem 3.4.

If $\alpha \neq 0$, then we may assume that $\alpha = -\frac{1}{2}$ and so $\mathsf{w} = \mathsf{w}_1 - 2\mathsf{w}_2 - 2\mathsf{w}_5$. Since

$$\partial_x \mathsf{w} = xy^2 + y^2x - 2yxy - 2x^3, \quad \partial_y \mathsf{w} = yx^2 + x^2y - 2xyx.$$

are linearly independent, $J(\mathsf{w})$ is standard by Proposition 3.3. It is easy to see that $\mathbf{M}(\mathsf{w}) = \begin{pmatrix} y^2-2x^2 & xy-2yx \\ yx-2xy & x^2 \end{pmatrix}$. Moreover we can calculate $\mathcal{V}(y^2-2x^2, xy-2yx, yx-2xy, x^2) = \emptyset$ in $\mathbb{P}^1\times\mathbb{P}^1$. Thus $J(\mathsf{w})$ is 3-dimensional Calabi-Yau by Theorem 3.4. The point scheme E of $J(\mathsf{w})$ is given by

$$\begin{aligned}\mathcal{V}(H(\mathsf{w})) &= \mathcal{V}\left(\det\begin{pmatrix} y_1y_2-2x_1x_2 & x_1y_2-2y_1x_2 \\ y_1x_2-2x_1y_2 & x_1x_2 \end{pmatrix}\right) \\ &= \mathcal{V}((x_1y_2-y_1x_2)^2 - x_1^2x_2^2) \\ &= \mathcal{V}(x_1y_2-y_1x_2-x_1x_2) \cup \mathcal{V}(x_1y_2-y_1x_2+x_1x_2) \subset \mathbb{P}^1\times\mathbb{P}^1\end{aligned}$$

	w (superpotential)	$\partial_x\mathsf{w}, \partial_y\mathsf{w}$ (defining relations of $J(\mathsf{w})$)	$H(\mathsf{w})$ (defining equation of E)	(E,τ), (geometric pair)
(1)	$\mathsf{w}_1 - 2\mathsf{w}_2$	$xy^2+y^2x-2yxy$, $yx^2+x^2y-2xyx$	$(x_1y_2-y_1x_2)^2$	$E = 2(1,1)$
(2)	$\mathsf{w}_1 - 2\mathsf{w}_2 - 2\mathsf{w}_5$	$xy^2+y^2x-2yxy-2x^3$, $yx^2+x^2y-2xyx$	$(x_1y_2-y_1x_2-x_1x_2)$ $\times(x_1y_2-y_1x_2+x_1x_2)$	$E=(1,1)+(1,1)$ meeting at 1 pt, τ stabilizes two components.
(3)	$\mathsf{w}_1 - 2\mathsf{w}_2 + \mathsf{w}_3$	$xy^2+y^2x-2yxy$ $+x^2y+xyx+yx^2$, $yx^2+x^2y-2xyx+x^3$	$2(x_1y_2-y_1x_2)^2$ $+2(y_1x_2+x_1y_2)x_1x_2$ $-x_1^2x_2^2$	E is an irr. curve with a cusp.
(4.1)	w_1	xy^2+y^2x, yx^2+x^2y	0	$E=\mathbb{P}^1\times\mathbb{P}^1$
(4.2)	$\mathsf{w}_1 + 2\mathsf{w}_2$	$xy^2+y^2x+2yxy$, $yx^2+x^2y+2xyx$	$(x_1y_2+y_1x_2)^2$	$E = 2(1,1)$
(4.3)	$\alpha\mathsf{w}_1+\beta\mathsf{w}_2$ $(\alpha\neq0,\beta\neq0,$ $\beta\neq\pm2\alpha)$	$\alpha(xy^2+y^2x)+\beta yxy$, $\alpha(yx^2+x^2y)+\beta xyx$	$\alpha x_1^2y_2^2+\alpha y_1^2x_2^2$ $+\beta x_1y_1x_2y_2$	$E=(1,1)+(1,1)$ meeting at 2 pts, τ stabilizes two components.
(5.1)	$\mathsf{w}_1 + 4\mathsf{w}_5$	$xy^2+y^2x+4x^3$, yx^2+x^2y	$x_1^2x_2^2$	$E=2(1,0)+2(0,1)$, τ interchanges two components
(5.2)	$\mathsf{w}_1 + 2\mathsf{w}_2 + 8\mathsf{w}_5$	$xy^2+y^2x+2yxy+8x^3$, $yx^2+x^2y+2xyx$	$(x_1y_2+y_1x_2-2x_1x_2)$ $\times(x_1y_2+y_1x_2+2x_1x_2)$	$E=(1,1)+(1,1)$ meeting at 1 pt, τ interchanges two components.
(5.3)	$\alpha\mathsf{w}_1+\beta\mathsf{w}_2+\mathsf{w}_5$ $(\alpha\neq0,\beta\neq0,$ $\beta\neq\pm2\alpha)$	$\alpha(xy^2+y^2x)+\beta yxy+x^3$, $\alpha(yx^2+x^2y)+\beta xyx$	$\alpha\beta x_1^2y_2^2+\alpha\beta y_1^2x_2^2$ $+\beta^2x_1y_1x_2y_2-\alpha x_1^2x_2^2$	E is an irr. curve with a biflecnode.
(6.1)	$\alpha\mathsf{w}_1+\beta\mathsf{w}_2+\mathsf{w}_5+\mathsf{w}_6$ $(\alpha=0,\beta\neq0,\pm1)$ or $(\beta=\pm1,\alpha\neq0,\pm1)$	$\gamma yxy+x^3$, $\gamma xyx+y^3$ $(\gamma\neq0,\pm1)$	$x_1y_1x_2y_2$	$E=(1,0)+(1,0)$ $+(0,1)+(0,1)$, τ circulates four components.
(6.2)	$\alpha\mathsf{w}_1+\beta\mathsf{w}_2+\mathsf{w}_5+\mathsf{w}_6$ $(\beta=0,\alpha\neq0,\pm\frac{1}{2})$ or $(\beta=2\alpha\pm1,\beta\neq0,\pm1)$	$xy^2+y^2x+\gamma x^3$, $yx^2+x^2y+\gamma y^3$ $(\gamma\neq0,\pm2)$	$x_1^2x_2^2+y_1^2y_2^2$ $+\gamma x_1y_1x_2y_2$	$E=(1,1)+(1,1)$ meeting at 2 pts, τ interchanges two components.
(6.3)	$\alpha\mathsf{w}_1+\beta\mathsf{w}_2+\mathsf{w}_5+\mathsf{w}_6$ $(\alpha\neq0,\beta\neq0,\pm1,$ $\beta\neq2\alpha\pm1,$ $\beta\neq-2\alpha\pm1)$	$\alpha(xy^2+y^2x)+\beta yxy+x^3$, $\alpha(yx^2+x^2y)+\beta xyx+y^3$	$\alpha\beta x_1^2y_2^2+\alpha\beta y_1^2x_2^2$ $+(\beta^2-1)x_1y_1x_2y_2$ $-\alpha x_1^2x_2^2-\alpha y_1^2y_2^2$	E is a smooth curve.

TABLE 1. Classification of Calabi-Yau superpotentials

(the union of two curves of bidegree (1, 1) meeting at one point), and the automorphism $\tau \in \mathrm{Aut}_k E$ is given by

$$\tau|_{\mathcal{V}(x_1y_2-y_1x_2-x_1x_2)}((p_1,q_1),(p_1,p_1+q_1)) = ((p_1,p_1+q_1),(p_1,2p_1+q_1)),$$
$$\tau|_{\mathcal{V}(x_1y_2-y_1x_2+x_1x_2)}((p_1,q_1),(p_1,-p_1+q_1)) = ((p_1,-p_1+q_1),(p_1,-2p_1+q_1))$$

(τ stabilizes two components).

(Case 3) If $\overline{\mathsf{w}} = 4f_3 = 4x^3y$, then $\mathsf{w} = \alpha\mathsf{w}_1 + \beta\mathsf{w}_2 + \mathsf{w}_3$ for $\alpha, \beta \in k$ such that $4\alpha + 2\beta = 0$. Since

$$\partial_x\mathsf{w} = \alpha(xy^2 + y^2x - 2yxy) + x^2y + xyx + yx^2, \quad \partial_y\mathsf{w} = \alpha(yx^2 + x^2y - 2xyx) + x^3$$

are linearly independent, $J(\mathsf{w})$ is standard by Proposition 3.3. It is easy to see that $\mathbf{M}(\mathsf{w}) = \begin{pmatrix} \alpha xy+yx+y^2 & \alpha(xy-2yx)+x^2 \\ \alpha(yx-2xy)+x^2 & \alpha x^2 \end{pmatrix}$. By Theorem 3.4, we can calculate that

$$\begin{aligned} &J(\mathsf{w}) \text{ is 3-dimensional Calabi-Yau} \\ &\iff \mathcal{V}(\alpha xy + yx + y^2, \alpha(xy - 2yx) + x^2, \alpha(yx - 2xy) + x^2, \alpha x^2) = \emptyset \text{ in } \mathbb{P}^1 \times \mathbb{P}^1 \\ &\iff \alpha \neq 0. \end{aligned}$$

Now assume that $\alpha \neq 0$. Then α may be equal to 1. The point scheme E of $J(\mathsf{w})$ is given by

$$\begin{aligned} \mathcal{V}(H(\mathsf{w})) =& \mathcal{V}\left(\det \begin{pmatrix} x_1y_2 + y_1x_2 + y_1y_2 & x_1y_2 - 2y_1x_2 + x_1x_2 \\ y_1x_2 - 2x_1y_2 + x_1x_2 & x_1x_2 \end{pmatrix}\right) \\ =& \mathcal{V}(2(x_1y_2 - y_1x_2)^2 + 2(y_1x_2 + x_1y_2)x_1x_2 - x_1^2x_2^2) \subset \mathbb{P}^1 \times \mathbb{P}^1 \end{aligned}$$

(an irreducible curve of bidegree (2, 2) with a cusp).

(Case 4) If $\overline{\mathsf{w}} = f_4 = x^2y^2$, then $\mathsf{w} = \alpha\mathsf{w}_1 + \beta\mathsf{w}_2$ for $\alpha, \beta \in k$ such that $4\alpha + 2\beta = 1$. Since

$$\partial_x\mathsf{w} = \alpha(xy^2 + y^2x) + \beta yxy, \quad \partial_y\mathsf{w} = \alpha(yx^2 + x^2y) + \beta xyx$$

are linearly independent, $J(\mathsf{w})$ is standard by Proposition 3.3. It is easy to see that $\mathbf{M}(\mathsf{w}) = \begin{pmatrix} \alpha y^2 & \alpha xy+\beta yx \\ \alpha yx+\beta xy & \alpha x^2 \end{pmatrix}$. By Theorem 3.4, we can calculate that

$$\begin{aligned} &J(\mathsf{w}) \text{ is 3-dimensional Calabi-Yau} \\ &\iff \mathcal{V}(\alpha y^2, \alpha xy + \beta yx, \alpha yx + \beta xy, \alpha x^2) = \emptyset \text{ in } \mathbb{P}^1 \times \mathbb{P}^1 \\ &\iff \alpha \neq 0. \end{aligned}$$

Now assume that $\alpha \neq 0$. Then the point scheme E of $J(\mathsf{w})$ is

$$\begin{aligned} \mathcal{V}(H(\mathsf{w})) =& \mathcal{V}\left(\det \begin{pmatrix} \alpha y_1y_2 & \alpha x_1y_2 + \beta y_1x_2 \\ \alpha y_1x_2 + \beta x_1y_2 & \alpha x_1x_2 \end{pmatrix}\right) \\ =& \mathcal{V}(\beta^2x_1y_1x_2y_2 + \alpha\beta x_1^2y_2^2 + \alpha\beta y_1^2x_2^2) \subset \mathbb{P}^1 \times \mathbb{P}^1. \end{aligned}$$

(4.1) If $\beta = 0$ (in this case $\alpha = \frac{1}{4}$), then $E = \mathbb{P}^1 \times \mathbb{P}^1$, and the automorphism $\tau \in \operatorname{Aut}_k E$ is given by

$$\tau((p_1, q_1), (p_2, q_2)) = ((p_2, q_2), (-p_1, q_1)).$$

(4.2) If $\beta \neq 0, \beta^2 - 4\alpha^2 = 0$ (in this case $\beta = 2\alpha$, so $\alpha = \frac{1}{8}, \beta = \frac{1}{4}$), then $E = \mathcal{V}((x_1y_2 + y_1x_2)^2)$ (a double curve of bidegree (1, 1)), and the automorphism $\tau \in \operatorname{Aut}_k E_{\mathrm{red}}$ is given by

$$\tau((p_1, q_1), (p_1, -q_1)) = ((p_1, -q_1), (p_1, q_1)).$$

(4.3) If $\beta \neq 0, \beta^2 - 4\alpha^2 \neq 0$ (that is, $\beta \neq 0, \beta \neq \pm 2\alpha$), then $E = \mathcal{V}(x_1y_2 - \gamma_1y_1x_2) \cup \mathcal{V}(x_1y_2 - \gamma_2y_1x_2)$ (the union of two curves of bidegree (1, 1) meeting at two points) where $\gamma_1 = \frac{-\beta + \sqrt{\beta^2 - 4\alpha^2}}{2\alpha}, \gamma_2 = \frac{-\beta - \sqrt{\beta^2 - 4\alpha^2}}{2\alpha}$, and the automorphism $\tau \in \operatorname{Aut}_k E$ is given by

$$\begin{aligned} \tau|_{\mathcal{V}(x_1y_2 - \gamma_1y_1x_2)}((p_1, q_1), (p_1, \gamma_1q_1)) &= ((p_1, \gamma_1q_1), (p_1, \gamma_1^2q_1)), \\ \tau|_{\mathcal{V}(x_1y_2 - \gamma_2y_1x_2)}((p_1, q_1), (p_1, \gamma_2q_1)) &= ((p_1, \gamma_2q_1), (p_1, \gamma_2^2q_1)) \end{aligned}$$

(τ stabilizes the components).

(Case 5) If $\overline{\mathsf{w}} = f_5 = x^4 + x^2y^2$, then $\mathsf{w} = \alpha\mathsf{w}_1 + \beta\mathsf{w}_2 + \mathsf{w}_5$ for $\alpha, \beta \in k$ such that $4\alpha + 2\beta = 1$. Since

$$\partial_x\mathsf{w} = \alpha(xy^2 + y^2x) + \beta yxy + x^3, \quad \partial_y\mathsf{w} = \alpha(yx^2 + x^2y) + \beta xyx$$

are linearly independent, $J(\mathsf{w})$ is standard by Proposition 3.3. It is easy to see that $\mathbf{M}(\mathsf{w}) = \begin{pmatrix} \alpha y^2+x^2 & \alpha xy+\beta yx \\ \alpha yx+\beta xy & \alpha x^2 \end{pmatrix}$. By Theorem 3.4, we can calculate that

$$\begin{aligned} &J(\mathsf{w}) \text{ is 3-dimensional Calabi-Yau} \\ \iff &\mathcal{V}(\alpha y^2 + x^2, \alpha xy + \beta yx, \alpha yx + \beta xy, \alpha x^2) = \emptyset \text{ in } \mathbb{P}^1 \times \mathbb{P}^1 \\ \iff &\alpha \neq 0. \end{aligned}$$

Now assume that $\alpha \neq 0$. Then the point scheme E of $J(\mathsf{w})$ is

$$\begin{aligned} \mathcal{V}(H(\mathsf{w})) =& \mathcal{V}\left(\det \begin{pmatrix} \alpha y_1y_2 + x_1x_2 & \alpha x_1y_2 + \beta y_1x_2 \\ \alpha y_1x_2 + \beta x_1y_2 & \alpha x_1x_2 \end{pmatrix}\right) \\ =& \mathcal{V}(\beta^2 x_1y_1x_2y_2 + \alpha\beta x_1^2y_2^2 + \alpha\beta y_1^2x_2^2 - \alpha x_1^2x_2^2) \subset \mathbb{P}^1 \times \mathbb{P}^1. \end{aligned}$$

(5.1) If $\beta = 0$ (in this case $\alpha = \frac{1}{4}$), then $E = \mathcal{V}(x_1^2) \cup \mathcal{V}(x_2^2)$ (the union of a double curve of bidegree (1, 0) and a double curve of bidegree (0, 1)), and the automorphism $\tau \in \mathrm{Aut}_k E_{\mathrm{red}}$ is given by

$$\begin{aligned} \tau|_{\mathcal{V}(x_1^2)}((0,1),(p_2,q_2)) &= ((p_2,q_2),(0,1)), \\ \tau|_{\mathcal{V}(x_2^2)}((p_1,q_1),(0,1)) &= ((0,1),(p_1,-q_1)) \end{aligned}$$

(σ interchanges two components).

(5.2) If $\beta \neq 0$, $\beta^2 - 4\alpha^2 = 0$ (in this case $\beta = 2\alpha$, so $\alpha = \frac{1}{8}, \beta = \frac{1}{4}$), then

$$E = \mathcal{V}(\frac{1}{2}(x_1y_2 + y_1x_2) - x_1x_2) \cup \mathcal{V}(\frac{1}{2}(x_1y_2 + y_1x_2) + x_1x_2)$$

(the union of two curves of bidegree (1, 1) meeting at one point), and the automorphism $\tau \in \mathrm{Aut}_k E$ is given by

$$\begin{aligned} \tau|_{\mathcal{V}(\frac{1}{2}(x_1y_2+y_1x_2)-x_1x_2)}((p_1,q_1),(p_1,2p_1-q_1)) &= ((p_1,2p_1-q_1),(p_1,-4p_1+q_1)), \\ \tau|_{\mathcal{V}(\frac{1}{2}(x_1y_2+y_1x_2)+x_1x_2)}((p_1,q_1),(p_1,-2p_1-q_1)) &= ((p_1,-2p_1-q_1),(p_1,4p_1+q_1)) \end{aligned}$$

(σ interchanges two components).

(5.3) If $\beta \neq 0$, $\beta^2 - 4\alpha^2 \neq 0$ (that is, $\beta \neq 0, \beta \neq \pm 2\alpha$), then

$$E = \mathcal{V}(\beta(x_1y_2 - \gamma_1y_1x_2)(x_1y_2 - \gamma_2y_1x_2) - \alpha x_1^2x_2^2)$$

(an irreducible curve of bidegree (2, 2) with a biflecnode) where $\gamma_1 = \frac{-\beta+\sqrt{\beta^2-4\alpha^2}}{2\alpha}, \gamma_2 = \frac{-\beta-\sqrt{\beta^2-4\alpha^2}}{2\alpha}$.

(Case 6) If $\overline{\mathsf{w}} = g_\lambda = x^4 + y^4 + \lambda x^2y^2$, then $\mathsf{w} = \alpha\mathsf{w}_1 + \beta\mathsf{w}_2 + \mathsf{w}_5 + \mathsf{w}_6$ for $\alpha, \beta \in k$ such that $4\alpha + 2\beta = \lambda (\neq \pm 2)$. Since

$$\partial_x\mathsf{w} = \alpha(xy^2 + y^2x) + \beta yxy + x^3, \quad \partial_y\mathsf{w} = \alpha(yx^2 + x^2y) + \beta xyx + y^3$$

are linearly independent, $J(\mathsf{w})$ is standard by Proposition 3.3. It is easy to see that $\mathbf{M}(\mathsf{w}) = \begin{pmatrix} \alpha y^2+x^2 & \alpha xy+\beta yx \\ \alpha yx+\beta xy & \alpha x^2+y^2 \end{pmatrix}$. By Theorem 3.4, we can calculate that

$$\begin{aligned} &J(\mathsf{w}) \text{ is 3-dimensional Calabi-Yau} \\ \iff &\mathcal{V}(\alpha y^2 + x^2, \alpha xy + \beta yx, \alpha yx + \beta xy, \alpha x^2 + y^2) = \emptyset \text{ in } \mathbb{P}^1 \times \mathbb{P}^1 \\ \iff &\alpha^2 - \beta^2 \neq 0 \text{ or } \alpha(\alpha^2 - 1) \neq 0 \\ \iff &(\alpha, \beta) \neq (0,0), (\pm 1, \pm 1) \qquad (\text{since } \lambda \neq \pm 2, (\alpha, \beta) \text{ can not be } (\pm 1, \mp 1)). \end{aligned}$$

Now assume that $(\alpha,\beta)\neq(0,0),(\pm1,\pm1)$. Then the point scheme E of $J(\mathsf{w})$ is

$$\begin{aligned}\mathcal{V}(H(\mathsf{w}))=&\mathcal{V}\left(\det\begin{pmatrix}\alpha y_1y_2+x_1x_2 & \alpha x_1y_2+\beta y_1x_2\\ \alpha y_1x_2+\beta x_1y_2 & \alpha x_1x_2+y_1y_2\end{pmatrix}\right)\\ =&\mathcal{V}((\beta^2-1)x_1y_1x_2y_2+\alpha\beta x_1^2y_2^2+\alpha\beta y_1^2x_2^2-\alpha x_1^2x_2^2-\alpha y_1^2y_2^2)\subset\mathbb{P}^1\times\mathbb{P}^1.\end{aligned}$$

If $\alpha=0,\beta^2-1=0$, then $\lambda=4\alpha+2\beta=\pm2$, so this case is excluded.

(6.1) If $\alpha=0,\beta^2-1\neq0$, then $E=\mathcal{V}(x_1)\cup\mathcal{V}(y_1)\cup\mathcal{V}(x_2)\cup\mathcal{V}(y_2)$ (the union of two curves of bidegree (1, 0) and two curves of bidegree (0, 1)), and the automorphism $\tau\in\operatorname{Aut}_k E$ is given by

$$\begin{aligned}\tau|_{\mathcal{V}(x_1)}((0,1),(p_2,q_2))&=((p_2,q_2),(1,0)),\\ \tau|_{\mathcal{V}(y_1)}((1,0),(p_2,q_2))&=((p_2,q_2),(0,1)),\\ \tau|_{\mathcal{V}(x_2)}((p_1,q_1),(0,1))&=((0,1),(q_1,-p_1)),\\ \tau|_{\mathcal{V}(y_2)}((p_1,q_1),(1,0))&=((1,0),(q_1,-p_1))\end{aligned}$$

(τ circulates four components).

If $\alpha\neq0,\beta=0,4\alpha^2-1=0$, then $\lambda=4\alpha+2\beta=\pm2$, so this case is excluded.

(6.2) If $\alpha\neq0,\beta=0,4\alpha^2-1\neq0$, then $E=\mathcal{V}(x_1x_2-\gamma_1y_1y_2)\cup\mathcal{V}(x_1x_2-\gamma_2y_1y_2)$ (the union of two curves of bidegree (1, 1) meeting at two points) where $\gamma_1=\frac{-1+\sqrt{1-4\alpha^2}}{2\alpha},\gamma_2=\frac{-1-\sqrt{1-4\alpha^2}}{2\alpha}$, and the automorphism $\tau\in\operatorname{Aut}_k E$ is given by

$$\begin{aligned}\tau|_{\mathcal{V}(x_1x_2-\gamma_1y_1y_2)}((p_1,q_1),(\gamma_1q_1,p_1))&=((\gamma_1q_1,p_1),(\gamma_2p_1,\gamma_1q_1)),\\ \tau|_{\mathcal{V}(x_1x_2-\gamma_2y_1y_2)}((p_1,q_1),(\gamma_2q_1,p_1))&=((\gamma_1q_1,p_1),(\gamma_1p_1,\gamma_2q_1))\end{aligned}$$

(τ interchanges two components).

We now consider the case $\alpha\neq0,\beta^2-1=0$. Since we can check that the case $\beta=-1$ is integrated with the case $\beta=1$, we may assume that $\beta=1$ (in this case $\alpha\neq\pm1$). Then

$$E=\mathcal{V}(x_1-y_1)\cup\mathcal{V}(x_1+y_1)\cup\mathcal{V}(x_2-y_2)\cup\mathcal{V}(x_2-y_2)$$

(the union of two curves of bidegree (1, 0) and two curves of bidegree (0, 1)), and the automorphism $\tau\in\operatorname{Aut}_k E$ is given by

$$\begin{aligned}\tau|_{\mathcal{V}(x_1-y_1)}((1,1),(p_2,q_2))&=((p_2,q_2),(1,-1)),\\ \tau|_{\mathcal{V}(x_1+y_1)}((1,-1),(p_2,q_2))&=((p_2,q_2),(1,1)),\\ \tau|_{\mathcal{V}(x_2-y_2)}((p_1,q_1),(1,1))&=((1,1),(\alpha p_1+q_1,-p_1-\alpha q_1)),\\ \tau|_{\mathcal{V}(x_2-y_2)}((p_1,q_1),(1,-1))&=((1,-1),(-\alpha p_1+q_1,-p_1+\alpha q_1))\end{aligned}$$

(τ circulates four components). Now $J(\mathsf{w})$ is $k\langle x,y\rangle(\alpha(xy^2+y^2x)+yxy+x^3,\alpha(yx^2+x^2y)+xyx+y^3)$. The homomorphism defined by $x\mapsto x-y,y\mapsto x+y$ induces the isomorphism

$$J(\mathsf{w})\xrightarrow{\sim}k\langle x,y\rangle/(\frac{1-\alpha}{1+\alpha}yxy+x^3,\frac{1-\alpha}{1+\alpha}xyx+y^3).$$

We can check that $\frac{1-\alpha}{1+\alpha}\neq0,\pm1$, so this case is integrated with the case (5.1).

We next consider the case $\alpha\neq0,\beta\neq0,\beta^2-1\neq0,2\alpha-\beta=\pm1$. Since we can check that the case $2\alpha-\beta=-1$ is integrated with the case $2\alpha-\beta=1$, we may assume that $2\alpha-\beta=1$. Then

$$E=\mathcal{V}(\sqrt{\beta}(x_1y_2+y_1x_2)+(x_1x_2+y_1y_2))\cup\mathcal{V}(\sqrt{\beta}(x_1y_2+y_1x_2)-(x_1x_2+y_1y_2))$$

(the union of two curves of bidegree (1, 1) meeting at two points), and the automorphism $\tau \in \mathrm{Aut}_k E$ is given by

$$\begin{aligned}
&\tau|_{\mathcal{V}(\sqrt{\beta}(x_1y_2+y_1x_2)+(x_1x_2+y_1y_2)}((p_1,q_1),(\sqrt{\beta}p_1+q_1,-p_1-\sqrt{\beta}q_1))\\
&=((\sqrt{\beta}p_1+q_1,-p_1-\sqrt{\beta}q_1)),((\beta+1)p_1+2\sqrt{\beta}q_1,2\sqrt{\beta}p_1+(\beta+1)q_1)),\\
&\tau|_{\mathcal{V}(\sqrt{\beta}(x_1y_2+y_1x_2)-(x_1x_2+y_1y_2)}((p_1,q_1),(\sqrt{\beta}p_1-q_1,p_1-\sqrt{\beta}q_1))\\
&=((\sqrt{\beta}p_1-q_1,p_1-\sqrt{\beta}q_1),((\beta+1)p_1-2\sqrt{\beta}q_1,-2\sqrt{\beta}p_1+(\beta+1)q_1))
\end{aligned}$$

(σ interchanges two components). Now $J(\mathsf{w})$ is $k\langle x,y\rangle(\alpha(xy^2+y^2x)+(2\alpha-1)yxy+x^3,\alpha(yx^2+x^2y)+(2\alpha-1)xyx+y^3)$. The homomorphism defined by $x\mapsto x-y, y\mapsto x+y$ induces the isomorphism

$$J(\mathsf{w})\xrightarrow{\sim} k\langle x,y\rangle/(\frac{1-\alpha}{2\alpha}(xy^2+y^2x)+x^3,\frac{1-\alpha}{2\alpha}(xy^2+y^2x)+y^3).$$

We can check that $\frac{1-\alpha}{2\alpha}\neq 0,\pm\frac{1}{2}$, so this case is integrated with the case (5.2).

(6.3) If $\alpha\neq 0,\beta\neq 0,\beta^2-1\neq 0,2\alpha-\beta\neq\pm 1$, then we can calculate that E is a smooth curve of bidegree (2, 2) by using the Jacobian criterion. □

Theorem 3.7. *Let V be a 2-dimensional vector space and $0\neq\mathsf{w}\in V^{\otimes 4}$. If $J(\mathsf{w})$ is not 3-dimensional Calabi-Yau, then $J(\mathsf{w})$ is isomorphic to one of the following five algebras:*

(1) $k\langle x,y\rangle/(x^3)$;
(2) $k\langle x,y\rangle/(x^3,x^2y+xyx+yx^2)$;
(3) $k\langle x,y\rangle/(yxy,xyx)$;
(4) $k\langle x,y\rangle/(yxy+x^3,xyx)$;
(5) $k\langle x,y\rangle/(x^3,y^3)$.

Proof. In each (Case 2), (Case 3), (Case 4), (Case 5) in the proof of Theorem 3.6, non-Calabi-Yau algebras are given by (1), (2), (3), (4) as in the above theorem. In (Case 6), non-Calabi-Yau algebras are isomorphic to either $k\langle x,y\rangle/(x^3,y^3)$ or $k\langle x,y\rangle/(xy^2+y^2x+yxy\pm x^3,yx^2+x^2y+xyx\pm y^3)$ by the proof of Theorem 3.6. For $a\in k$,

$$(x+ay)^3=a^2(xy^2+y^2x+yxy)+x^3+a(yx^2+x^2y+xyx)+a^3y^3,$$

so we have

$$\begin{aligned}
&k\langle x,y\rangle/(xy^2+y^2x+yxy+x^3,yx^2+x^2y+xyx+y^3)=k\langle x,y\rangle/((x+y)^3,(x-y)^3),\\
&k\langle x,y\rangle/(xy^2+y^2x+yxy-x^3,yx^2+x^2y+xyx-y^3)=k\langle x,y\rangle/((x+\sqrt{-1}y)^3,(x-\sqrt{-1}y)^3),
\end{aligned}$$

hence the result. □

Corollary 3.8. *Let V be a 2-dimensional vector space and $0\neq\mathsf{w}\in V^{\otimes 4}$. Then $J(\mathsf{w})$ is 3-dimensional Calabi-Yau if and only if it is a domain.*

Proof. If $J(\mathsf{w})$ is 3-dimensional Calabi-Yau, then it is 3-dimensional noetherian AS-regular, so it is a domain by [3, Theorem 3.9]. On the other hand, if $J(\mathsf{w})$ is not 3-dimensional Calabi-Yau, then it is not a domain by the classification in Theorem 3.7. □

4. Homological Determinants

The homological determinant plays an important role in invariant theory for AS-regular algebras ([8], [10], [11], [6], [14] etc.). For a 3-dimensional noetherian quadratic Calabi-Yau algebra $S=J(\mathsf{w})=T(V)/(R)$, it was shown in [12, Theorem 7.2] that $\mathrm{hdet}\,\sigma=\det\sigma|_V$ for every $\sigma\in\mathrm{GrAut}\,S$ if and only if $c(\mathsf{w})\notin\mathrm{Sym}^3 V$. In this section, we will compute homological determinants for 3-dimensional noetherian cubic Calabi-Yau algebras.

Lemma 4.1. *Let $n = \dim V$, $m \in \mathbb{N}^+$, and $\sigma \in \mathrm{GL}(V)$. For $\mathsf{w} \in (\mathrm{Alt}^n V)^{\otimes m}$, $\sigma(\mathsf{w}) = (\det \sigma|_V)^m \mathsf{w}$.*

Proof. Fix a basis $x_1, \dots, x_n$ for V. First, we show it for $m = 1$. Let $\mathsf{w} = x_{i_1} \otimes \cdots \otimes x_{i_n} \in V^{\otimes n}$ be a monomial. Since $a(\theta(\mathsf{w})) = \mathrm{sgn}(\theta) a(\mathsf{w})$ for $\theta \in S_n$, if $x_{i_s} = x_{i_t}$ for some $s \neq t$, then $a(\mathsf{w}) = 0$, and if $i_s \neq i_t$ for every $s \neq t$, then $a(\mathsf{w}) = \pm a(x_1 \otimes \cdots \otimes x_n)$, so $\mathrm{Alt}^n V = \mathrm{Im}\, a$ is a one dimensional vector space spanned by $\mathsf{w}_0 := a(x_1 \otimes \cdots \otimes x_n)$. Since $S = \mathcal{D}(\mathsf{w}_0, n-2) = k[x_1, \dots, x_n]$ is the polynomial algebra, σ extends to a grade algebra automorphism of S, so $\sigma(\mathsf{w}_0) = (\mathrm{hdet}\, \sigma)\mathsf{w}_0$ by [12, Theorem 3.3], but for a polynomial algebra, it is known that $\mathrm{hdet}\, \sigma = \det \sigma|_V$, hence $\sigma(\mathsf{w}_0) = (\det \sigma|_V)\mathsf{w}_0$.

For $m > 1$, if $\mathsf{w} = \mathsf{v}_1 \otimes \cdots \otimes \mathsf{v}_m \in (\mathrm{Alt}^n V)^{\otimes m}$ where $\mathsf{v}_i \in \mathrm{Alt}^n V$, then

$$\sigma(\mathsf{w}) = \sigma(\mathsf{v}_1) \otimes \cdots \otimes \sigma(\mathsf{v}_m) = (\det \sigma|_V)\mathsf{v}_1 \otimes \cdots \otimes (\det \sigma|_V)\mathsf{v}_m = (\det \sigma|_V)^m \mathsf{w}.$$

The result follows by linearity. □

Let V be a 2-dimensional vector space. Since $V^{\otimes 2} = \mathrm{Sym}^2 V \oplus \mathrm{Alt}^2 V$,

$$\begin{aligned} V^{\otimes 4} &= (\mathrm{Sym}^2 V \oplus \mathrm{Alt}^2 V) \otimes (\mathrm{Sym}^2 V \oplus \mathrm{Alt}^2 V) \\ &= (\mathrm{Sym}^2 V)^{\otimes 2} \oplus (\mathrm{Sym}^2 V \otimes \mathrm{Alt}^2 V) \oplus (\mathrm{Alt}^2 V \otimes \mathrm{Sym}^2 V) \oplus (\mathrm{Alt}^2 V)^{\otimes 2}. \end{aligned}$$

Let $\pi : V^{\otimes 4} \to (\mathrm{Alt}^2 V)^{\otimes 2}$ be the projection map with respect to the above decomposition. Fix a basis x, y for V. Since $\mathrm{Sym}^2 V = kx^2 + k(xy + yx) + ky^2$ and $\mathrm{Alt}^2 V = k(xy - yx)$, we have

$$\begin{aligned} (\mathrm{Sym}^2 V)^{\otimes 2} &= kx^4 + ky^4 + kx^2y^2 + ky^2x^2 + k(x^3y + x^2yx) + k(xyx^2 + yx^3) \\ &\quad + k(y^2xy + y^3x) + k(xy^3 + yxy^2) + k(xyxy + xy^2x + yx^2y + yxyx), \\ \mathrm{Sym}^2 V \otimes \mathrm{Alt}^2 V &= k(x^3y - x^2yx) + k(y^2xy - y^3x) + k(xyxy - xy^2x + yx^2y - yxyx), \\ \mathrm{Alt}^2 V \otimes \mathrm{Sym}^2 V &= k(xyx^2 - yx^3) + k(xy^3 - yxy^2) + k(xyxy + xy^2x - yx^2y - yxyx), \\ (\mathrm{Alt}^2 V)^{\otimes 2} &= k(xyxy - xy^2x - yx^2y + yxyx). \end{aligned}$$

Let $\mathsf{w}_0 = xyxy - xy^2x - yx^2y + yxyx$ so that $(\mathrm{Alt}^2 V)^{\otimes 2} = k\mathsf{w}_0$, and define a map $\mu : V^{\otimes 4} \to k$ by $\pi(\mathsf{w}) = \mu(\mathsf{w})\mathsf{w}_0$.

Remark 4.2. Although the map μ depends on the choice of a basis for V, the map π is independent of the choice of a basis for V. Since $\mu(\mathsf{w}) = 0$ if and only if $\pi(\mathsf{w}) = 0$, whether $\mu(\mathsf{w}) = 0$ or not is independent of the choice of a basis for V.

Theorem 4.3. *Let $S = T(V)/(R)$ be a 3-dimensional noetherian cubic AS-regular algebra where $R \subset V^{\otimes 3}$. If $\mu(\mathsf{w}_S) \neq 0$, then $\mathrm{hdet}\, \sigma = (\det \sigma|_V)^2$ for every $\sigma \in \mathrm{GrAut}\, S$.*

Proof. Write $W_s := (\mathrm{Sym}^2 V)^{\otimes 2} \oplus (\mathrm{Sym}^2 V \otimes \mathrm{Alt}^2 V) \oplus (\mathrm{Alt}^2 V \otimes \mathrm{Sym}^2 V)$ and $W_a := (\mathrm{Alt}^2 V)^{\otimes 2}$ so that $V^{\otimes 4} = W_s \oplus W_a$. For $\mathsf{w} \in V^{\otimes 2}$, $\sigma(s(\mathsf{w})) = s(\sigma(\mathsf{w}))$ and $\sigma(a(\mathsf{w})) = a(\sigma(\mathsf{w}))$ by Lemma 2.3, so $\sigma(W_s) = W_s$ and $\sigma(W_a) = W_a$.

We may write $\mathsf{w}_S = \mathsf{w}_s + \mathsf{w}_a$ where $\mathsf{w}_a := \pi(\mathsf{w}_S) \in W_a, \mathsf{w}_s := \mathsf{w}_S - \pi(\mathsf{w}_S) \in W_s$ in a unique way. By [12, Theorem 3.3] and Lemma 4.1,

$$(\mathrm{hdet}\, \sigma)(\mathsf{w}_s + \mathsf{w}_a) = (\mathrm{hdet}\, \sigma)(\mathsf{w}_S) = \sigma(\mathsf{w}_S) = \sigma(\mathsf{w}_s + \mathsf{w}_a) = \sigma(\mathsf{w}_s) + (\det \sigma|_V)^2 \mathsf{w}_a,$$

so

$$(\mathrm{hdet}\, \sigma - (\det \sigma|_V)^2)\mathsf{w}_a = \sigma(\mathsf{w}_s) - (\mathrm{hdet}\, \sigma)\mathsf{w}_s = 0$$

because the left hand side is in W_a and the right hand side is in W_s. If $\mu(\mathsf{w}_S) \neq 0$, then $\mathsf{w}_a = \pi(\mathsf{w}_S) \neq 0$, so $\mathrm{hdet}\, \sigma = (\det \sigma|_V)^2$. □

Example 4.4. If $S = k\langle x, y\rangle/(x^2y - yx^2, y^2x - xy^2)$, then S is a 3-dimensional noetherian cubic AS-regular algebra (but not a Calabi-Yau algebra) such that

$$\begin{aligned}\mathsf{w}_S &= xy^2x - x^2y^2 + yx^2y - y^2x^2\\ &= -\frac{1}{2}(xyxy - xy^2x - yx^2y + yxyx) + \frac{1}{2}(xyxy + xy^2x + yx^2y + yxyx) - x^2y^2 - y^2x^2\\ &= -\frac{1}{2}\mathsf{w}_0 + \mathsf{w}'\end{aligned}$$

where $\mathsf{w}' \in (\mathrm{Sym}^2 V)^{\otimes 2}$, so $\mu(\mathsf{w}_S) = -\frac{1}{2} \neq 0$, hence $\mathrm{hdet}\,\sigma = (\det\sigma|_V)^2$ for every $\sigma \in \mathrm{GrAut}\,S$ by Theorem 4.3. In fact, it is easy to calculate $\sigma(\mathsf{w}_S)$ for elementary matrices $\sigma \in \mathrm{GL}(2,k)$:

σ	$\begin{pmatrix}0 & 1\\ 1 & 0\end{pmatrix}$	$\begin{pmatrix}\alpha & 0\\ 0 & 1\end{pmatrix}$	$\begin{pmatrix}1 & \alpha\\ 0 & 1\end{pmatrix}$
$\det(\sigma)$	-1	α	1
$\sigma(\mathsf{w}_S)$	w_S	$\alpha^2\mathsf{w}_S$	w_S
$\mathrm{hdet}(\sigma)$	1	α^2	1

Since every $\sigma \in \mathrm{GL}(2,k)$ is a product of elementary matrices, every σ extends to a graded algebra automorphism $\sigma \in \mathrm{GrAut}\,S$ by [12, Theorem 3.2]. Since both det and hdet are group homomorphisms, $\mathrm{hdet}\,\sigma = (\det\sigma|_V)^2$ for every $\sigma \in \mathrm{GrAut}\,S$ by the above table.

Recall that we set $W' := \bigoplus_{i=3}^{6} k\mathsf{w}_i \subset \mathrm{Sym}^4 V$ where w_i are as defined in Section 3.3.

Lemma 4.5. *Let V be a 2-dimensional vector space. For a superpotential $\mathsf{w} \in \mathrm{Im}\,c \subset V^{\otimes 4}$, $\mathsf{w} \in \mathrm{Sym}^4 V$ if and only if $\mu(\mathsf{w}) = 0$.*

Proof. Every $\mathsf{w} \in \mathrm{Im}\,c$ can be written as $\mathsf{w} = \alpha(\mathsf{w}_2 - \mathsf{w}_1) + \beta(\mathsf{w}_2 + \mathsf{w}_1) + \mathsf{w}'$ where $\alpha, \beta \in k$ and $\mathsf{w}' \in W'$. It is easy to see that $\mathsf{w}_2 + \mathsf{w}_1, \mathsf{w}' \in (\mathrm{Sym}^2 V)^{\otimes 2}$. Since $x^2y^2, y^2x^2 \in (\mathrm{Sym}^2 V)^{\otimes 2}$,

$$\mu(\mathsf{w}) = \alpha\mu(\mathsf{w}_2 - \mathsf{w}_1) = \alpha\mu(\mathsf{w}_0 + x^2y^2 + y^2x^2) = \alpha.$$

Since $\mathsf{w} = (\beta - \alpha)\mathsf{w}_1 + (\beta + \alpha)\mathsf{w}_2 + \mathsf{w}'$, $\mathsf{w} \in \mathrm{Sym}^4 V$ if and only if $\beta - \alpha = \beta + \alpha$ if and only if $\mu(\mathsf{w}) = \alpha = 0$. □

Corollary 4.6. *Let $S = T(V)/(R)$ be a 3-dimensional noetherian cubic Calabi-Yau algebra where $R \subset V^{\otimes 3}$. If $\mathsf{w}_S \notin \mathrm{Sym}^4 V$, then $\mathrm{hdet}\,\sigma = (\det\sigma|_V)^2$ for every $\sigma \in \mathrm{GrAut}\,S$.*

Proof. Since S is Calabi-Yau, $\mathsf{w}_S \in \mathrm{Im}\,c$ is a superpotential, so the result follows from Theorem 4.3 and Lemma 4.5. □

Proposition 4.7. *Every 3-dimensional noetherian cubic Calabi-Yau algebra S such that $\mathsf{w}_S \in \mathrm{Sym}^4 V$ is isomorphic to one of the following algebras:*

1. $k\langle x, y\rangle/(xy^2 + yxy + y^2x, yx^2 + xyx + x^2y)$;
2. $k\langle x, y\rangle/(xy^2 + yxy + y^2x + x^3, yx^2 + xyx + x^2y)$;
3. $k\langle x, y\rangle/(xy^2 + yxy + y^2x + ax^3, yx^2 + xyx + x^2y + ay^3)$, $a \in k \setminus \{0, \pm 1, \pm 3\}$.

Proof. This follows from the proof of Theorem 3.6. □

Theorem 4.8. *Let S be a 3-dimensional noetherian cubic Calabi-Yau algebra. Then $\mathrm{hdet}\,\sigma = (\det\sigma|_V)^2$ for every $\sigma \in \mathrm{GrAut}\,S$ if and only if*

$$S \ncong k\langle x, y\rangle/(xy^2 + yxy + y^2x + \sqrt{-3}x^3, yx^2 + xyx + x^2y + \sqrt{-3}y^3).$$

Proof. By Corollary 4.6, it is enough to check the algebras in Proposition 4.7. Let $\mathsf{w} \in \operatorname{Im} c \subset V^{\otimes 4}$ be a Calabi-Yau superpotential. Then $\sigma \in \operatorname{GL}(V)$ extends to $\sigma \in \operatorname{GrAut} J(\mathsf{w})$ if and only if $\sigma(\mathsf{w}) = \lambda \mathsf{w}$ for some $\lambda \in k$ by [12, Theorem 3.2]. In this case, $\lambda = \operatorname{hdet} \sigma$ by [12, Theorem 3.3], and $\sigma(\overline{\mathsf{w}}) = \overline{\sigma(\mathsf{w})} = \overline{\lambda \mathsf{w}} = \lambda \overline{\mathsf{w}}$ by Lemma 2.5.

(Case 1) If $S = k\langle x, y\rangle/(xy^2 + yxy + y^2x, yx^2 + xyx + x^2y)$, then $\mathsf{w}_S = x^2y^2 + xy^2x + y^2x^2 + yx^2y + xyxy + yxyx \in \operatorname{Sym}^4 V$. Since $\overline{\mathsf{w}}_S = 6x^2y^2$, if $\sigma = \left(\begin{smallmatrix} \alpha & \beta \\ \gamma & \delta \end{smallmatrix}\right) \in \operatorname{GL}(2,k)$ extends to $\sigma \in \operatorname{GrAut} S$, then

$$\sigma(x^2y^2) = (\alpha x + \beta y)^2(\gamma x + \delta y)^2 = \alpha^2\gamma^2 x^4 + \cdots + \beta^2\delta^2 y^4 = \lambda x^2y^2$$

for some $\lambda \in k$, so $\alpha\gamma = \beta\delta = 0$. Since $\alpha\delta - \beta\gamma \neq 0$, it follows that either $\sigma = \left(\begin{smallmatrix} \alpha & 0 \\ 0 & \delta \end{smallmatrix}\right)$ or $\sigma = \left(\begin{smallmatrix} 0 & \beta \\ \gamma & 0 \end{smallmatrix}\right)$. If $\sigma = \left(\begin{smallmatrix} \alpha & 0 \\ 0 & \delta \end{smallmatrix}\right)$, then $\operatorname{hdet} \sigma = \alpha^2\delta^2$ and $\det \sigma|_V = \alpha\delta$. If $\sigma = \left(\begin{smallmatrix} 0 & \beta \\ \gamma & 0 \end{smallmatrix}\right)$, then $\operatorname{hdet} \sigma = \beta^2\gamma^2$ and $\det \sigma|_V = -\beta\gamma$. In either case, $\operatorname{hdet} \sigma = (\det \sigma|_V)^2$.

(Case 2) If $S = k\langle x, y\rangle/(xy^2 + yxy + y^2x + x^3, yx^2 + xyx + x^2y)$, then $\mathsf{w}_S = x^2y^2 + xy^2x + y^2x^2 + yx^2y + xyxy + yxyx + x^4 \in \operatorname{Sym}^4 V$. Since $\overline{\mathsf{w}}_S = 6x^2y^2 + x^4$, if $\sigma = \left(\begin{smallmatrix} \alpha & \beta \\ \gamma & \delta \end{smallmatrix}\right) \in \operatorname{GL}(2,k)$ extends to $\sigma \in \operatorname{GrAut} S$, then $\sigma(6x^2y^2 + x^4) = \lambda(6x^2y^2 + x^4)$ for some $\lambda \in k$. Since

$$\begin{aligned}\sigma(6x^2y^2 + x^4) &= 6(\alpha x + \beta y)^2(\gamma x + \delta y)^2 + (\alpha x + \beta y)^4 \\ &= \alpha^2(6\gamma^2 + \alpha^2)x^4 + 4\alpha(3\alpha\gamma\delta + 3\beta\gamma^2 + \alpha^2\beta)x^3y + 6(\alpha^2\delta^2 + 4\alpha\beta\gamma\delta + \beta^2\gamma^2 + \alpha^2\beta^2)x^2y^2 \\ &\quad + 4\beta(3\beta\gamma\delta + 3\alpha\delta^2 + \alpha\beta^2)xy^3 + \beta^2(6\delta^2 + \beta^2)y^4,\end{aligned}$$

we obtain $4\alpha(3\alpha\gamma\delta + 3\beta\gamma^2 + \alpha^2\beta) = 0$, $4\beta(3\beta\gamma\delta + 3\alpha\delta^2 + \alpha\beta^2) = 0$, $\beta^2(6\delta^2 + \beta^2) = 0$, and $\alpha^2\delta^2 + 4\alpha\beta\gamma\delta + \beta^2\gamma^2 + \alpha^2\beta^2 = \alpha^2(6\gamma^2 + \alpha^2)$. If $\beta \neq 0$, then $\beta^2 = -6\delta^2$, so $0 = 4\beta(3\beta\gamma\delta + 3\alpha\delta^2 + \alpha\beta^2) = 4\beta(3\beta\gamma\delta - 3\alpha\delta^2) = -3\beta\delta(\alpha\delta - \beta\gamma)$. This is a contradiction, so $\beta = 0$. Then we have $\alpha^2\gamma\delta = 0$. Since $\alpha\delta - \beta\gamma = \alpha\delta \neq 0$, we see $\gamma = 0$. Therefore $\sigma = \left(\begin{smallmatrix} \alpha & 0 \\ 0 & \delta \end{smallmatrix}\right)$ and $\alpha^2 = \delta^2$. We can check that $\det \sigma|_V = \alpha\delta$ and $\operatorname{hdet} \sigma = \alpha^2\delta^2$, so $\operatorname{hdet} \sigma = (\det \sigma|_V)^2$.

(Case 3) If $S = k\langle x, y\rangle/(xy^2 + yxy + y^2x + ax^3, yx^2 + xyx + x^2y + ay^3)$ where $a \in k \setminus \{0, \pm 1, \pm 3\}$, then $\mathsf{w}_S = x^2y^2 + xy^2x + y^2x^2 + yx^2y + xyxy + yxyx + ax^4 + ay^4 \in \operatorname{Sym}^4 V$. Since $\overline{\mathsf{w}}_S = 6x^2y^2 + ax^4 + ay^4$, if $\sigma = \left(\begin{smallmatrix} \alpha & \beta \\ \gamma & \delta \end{smallmatrix}\right) \in \operatorname{GL}(2,k)$ extends to $\sigma \in \operatorname{GrAut} S$, then $\sigma(6x^2y^2 + ax^4 + ay^4) = \lambda(6x^2y^2 + ax^4 + ay^4)$ for some $\lambda \in k$. Since

$$\begin{aligned}\sigma(6x^2y^2 + ax^4 + ay^4) &= 6(\alpha x + \beta y)^2(\gamma x + \delta y)^2 + a(\alpha x + \beta y)^4 + a(\gamma x + \delta y)^4 \\ &= (6\alpha^2\gamma^2 + a\alpha^4 + a\gamma^4)x^4 + 4(3\alpha^2\gamma\delta + 3\alpha\beta\gamma^2 + a\alpha^3\beta + a\gamma^3\delta)x^3y \\ &\quad + 6(\alpha^2\delta^2 + 4\alpha\beta\gamma\delta + \beta^2\delta^2 + a\alpha^2\beta^2 + a\gamma^2\delta^2)x^2y^2 \\ &\quad + 4(3\beta^2\gamma\delta + 3\alpha\beta\delta^2 + a\alpha\beta^3 + a\gamma\delta^3)xy^3 + (6\beta^2\delta^2 + a\beta^4 + a\delta^4)y^4,\end{aligned}$$

it follows that

$$3\alpha^2\gamma\delta + 3\alpha\beta\gamma^2 + a\alpha^3\beta + a\gamma^3\delta = 0, \quad 3\beta^2\gamma\delta + 3\alpha\beta\delta^2 + a\alpha\beta^3 + a\gamma\delta^3 = 0, \text{ and} \tag{4.1}$$

$$\alpha^2\delta^2 + 4\alpha\beta\gamma\delta + \beta^2\delta^2 + a\alpha^2\beta^2 + a\gamma^2\delta^2 = a^{-1}(6\alpha^2\gamma^2 + a\alpha^4 + a\gamma^4) = a^{-1}(6\beta^2\delta^2 + a\beta^4 + a\delta^4). \tag{4.2}$$

By the equations in (4.1), we have $(3\alpha\beta + a\gamma\delta)(\alpha\delta + \beta\gamma)(\alpha\delta - \beta\gamma) = 0$.

Assume that $3\alpha\beta + a\gamma\delta = 0$. Then $(9 - a^2)\alpha^2\gamma\delta = (9 - a^2)\beta^2\gamma\delta = 0$ by (4.1). Since $a \neq \pm 3$ and $\alpha\delta - \beta\gamma \neq 0$, it follows that either $\gamma = 0$ or $\delta = 0$. If $\gamma = 0$, then $\beta = 0$, so $\sigma = \left(\begin{smallmatrix} \alpha & 0 \\ 0 & \delta \end{smallmatrix}\right)$ and $\alpha^2 = \delta^2$ by (4.2). We can check that $\det \sigma|_V = \alpha\delta$ and $\operatorname{hdet} \sigma = \alpha^2\delta^2$, so $\operatorname{hdet} \sigma = (\det \sigma|_V)^2$. If $\delta = 0$, then $\alpha = 0$, so $\sigma = \left(\begin{smallmatrix} 0 & \beta \\ \gamma & 0 \end{smallmatrix}\right)$ and $\beta^2 = \gamma^2$ by (4.2). We can check that $\det \sigma|_V = -\beta\gamma$ and $\operatorname{hdet} \sigma = \beta^2\gamma^2$, so $\operatorname{hdet} \sigma = (\det \sigma|_V)^2$.

Assume that $\alpha\delta+\beta\gamma=0$. Then $\alpha^4=\gamma^4, \beta^4=\delta^4$ by (4.1), so $\sigma=\left(\begin{smallmatrix}\alpha & \beta\\ -\xi\alpha & \xi\beta\end{smallmatrix}\right)$ where ξ is a 4th root of unity. By (4.2), we see $(3\xi^2+a)(\alpha^4+\beta^4)=0$. Since $a\neq\pm3$, it follows that $\beta^2=\pm\alpha^2$. Thus, by (4.2), we have $a^2+(1-\xi^2)a+3\xi^2=0$ or $a^2-(1+\xi^2)a-3\xi^2=0$. Since $\xi^2=\pm1$ and $a\neq\pm1,\pm3$, we see $a=\pm\sqrt{-3}$. Hence it follows that unless $a=\pm\sqrt{-3}$, $\operatorname{hdet}\sigma=(\det\sigma|_V)^2$ for every $\sigma\in\operatorname{GrAut}S$.

Conversely, suppose that $S=k\langle x,y\rangle/(xy^2+yxy+y^2x+\sqrt{-3}x^3, yx^2+xyx+x^2y+\sqrt{-3}y^3)$. If $\sigma=\left(\begin{smallmatrix}\alpha & \beta\\ -\xi\alpha & \xi\beta\end{smallmatrix}\right)$ where $\beta^2=\pm\alpha^2, \xi^2=\mp1$,then we can check that $\sigma\in\operatorname{GL}(2,k)$ extends to $\sigma\in\operatorname{GrAut}S$, and $\det\sigma|_V=2\xi\alpha\beta$, so $(\det\sigma|_V)^2=-4\alpha^4$. Since

$$\begin{aligned}
&\sigma(6x^2y^2+\sqrt{-3}x^4+\sqrt{-3}y^4)\\
&=\alpha^4(-2\sqrt{-3}\xi^2+2)\sqrt{-3}x^4+(-2\alpha^2\beta^2\xi^2+2\sqrt{-3}\alpha^2\beta^2)6x^2y^2+\beta^4(-2\sqrt{-3}\xi^2+2)\sqrt{-3}y^4\\
&=\alpha^4(2\pm2\sqrt{-3})\sqrt{-3}x^4+(2\alpha^4\pm2\sqrt{-3}\alpha^4)6x^2y^2+\alpha^4(2\pm2\sqrt{-3})\sqrt{-3}y^4\\
&=\alpha^4(2\pm2\sqrt{-3})(6x^2y^2+\sqrt{-3}x^4+\sqrt{-3}y^4),
\end{aligned}$$

we obtain $\operatorname{hdet}\sigma=\alpha^4(2\pm2\sqrt{-3})=-4\alpha^4(\frac{-1\mp\sqrt{-3}}{2})=-4\alpha^4\omega=(\det\sigma|_V)^2\omega\neq(\det\sigma|_V)^2$ where ω is a primitive 3rd root of unity.

The homomorphism defined by $x\mapsto\sqrt{-1}x, y\mapsto y$ induces the isomorphism

$$\begin{aligned}
&k\langle x,y\rangle/(xy^2+yxy+y^2x+\sqrt{-3}x^3, yx^2+xyx+x^2y+\sqrt{-3}y^3)\\
&\cong k\langle x,y\rangle/(xy^2+yxy+y^2x-\sqrt{-3}x^3, yx^2+xyx+x^2y-\sqrt{-3}y^3),
\end{aligned}$$

so we have the result. □

5. Connections to Other Classes of Algebras

Let $S=J(\mathsf{w})=T(V)/(R)$ be a 3-dimensional noetherian quadratic Calabi-Yau algebra. In [13], it was shown that $c(\mathsf{w})\notin\operatorname{Sym}^3V$ if and only if S is a deformation quantization of the polynomial algebra $k[x,y,z]$, and $c(\mathsf{w})\in\operatorname{Sym}^3V$ if and only if S is a Clifford algebra. In this last section, we will see something similar holds at least in one direction.

5.1. **Deformation Quantizations.** For $\lambda\in k$ and $f\in k[x,y,z]_3$ where $\deg x=\deg y=\deg z=1$, the algebra $S_f^\lambda:=k\langle x,y,z\rangle/([y,z]-\lambda\widetilde{f_x},[z,x]-\lambda\widetilde{f_y},[x,y]-\lambda\widetilde{f_z})$ is a deformation quantization of $k[x,y,z]$ ([7]), which is often a 3-dimensional noetherian quadratic Calabi-Yau algebra ([13]). For $\lambda\in k$ and $f\in k[x,y,z]_4$ where $\deg x=\deg y=1$, $\deg z=2$, we may still define $S_f^\lambda:=k\langle x,y,z\rangle/([y,z]-\lambda\widetilde{f_x},[z,x]-\lambda\widetilde{f_y},[x,y]-\lambda\widetilde{f_z})$. We do not know if S_f^λ is a deformation quantization of $k[x,y,z]$, but we will study when a 3-dimensional noetherian cubic Calabi-Yau algebra is of the form S_f^λ.

Theorem 5.1. *Let V be a 2-dimensional vector space and $\mathsf{w}\in V^{\otimes4}$. Fix a basis x,y for V. If $c(\mathsf{w})\notin\operatorname{Sym}^4V$, then $J(\mathsf{w})=S_f^\lambda$ where $\lambda=-\frac{3}{8\mu(c(\mathsf{w}))}\in k$ and $f=\overline{\mathsf{w}}+\frac{4\mu(c(\mathsf{w}))}{3}z^2\in k[x,y,z]_4$ with $\deg x=\deg y=1,\deg z=2$.*

Proof. Since $J(c(\mathsf{w}))=J(\mathsf{w})$, we may assume that $\mathsf{w}=c(\mathsf{w})\in\operatorname{Im}c$ is a superpotential. If V is a 2-dimensional vector space spanned by x,y, then we may write $\mathsf{w}=\alpha\mathsf{w}_1+\beta\mathsf{w}_2+\mathsf{w}'$ where $\alpha,\beta\in k$ and $\mathsf{w}'\in W'$, and, in this case,

$$\partial_x\mathsf{w}=\alpha(xy^2+y^2x)+\beta yxy+\partial_x\mathsf{w}',\quad \partial_y\mathsf{w}=\alpha(yx^2+x^2y)+\beta xyx+\partial_y\mathsf{w}'.$$

Note that $\mathsf{w}\notin\operatorname{Sym}^4V$ if and only if $\alpha\neq\beta$, and, in this case, we set $\lambda=\frac{3}{4(\alpha-\beta)}$ and $f=\overline{\mathsf{w}}-\frac{2(\alpha-\beta)}{3}z^2$. Since $f=2(2\alpha+\beta)x^2y^2+f'-\frac{2(\alpha-\beta)}{3}z^2$ where $f'=\overline{\mathsf{w}'}$, it follows that $[x,y]-\lambda\widetilde{f_z}=xy-yx-z$.

Moreover, by Lemma 2.6,

$$\begin{aligned}
[y,z]-\lambda\widetilde{f_x} &= yz-zy-\frac{3}{4(\alpha-\beta)}(4(2\alpha+\beta)\widetilde{xy^2}+\widetilde{f'_x})\\
&= y(xy-yx)-(xy-yx)y-\frac{3(2\alpha+\beta)}{\alpha-\beta}\frac{xy^2+yxy+y^2x}{3}-\frac{3}{4(\alpha-\beta)}\widetilde{f'_x}\\
&= -(\frac{2\alpha+\beta}{\alpha-\beta}+1)(xy^2+y^2x)-(\frac{2\alpha+\beta}{\alpha-\beta}-2)yxy-\frac{3}{4(\alpha-\beta)}4\partial_x\mathsf{w}'\\
&= -\frac{3}{\alpha-\beta}(\alpha(xy^2+y^2x)+\beta yxy+\partial_x\mathsf{w}')\\
&= -\frac{3}{\alpha-\beta}\partial_x\mathsf{w}
\end{aligned}$$

and, similarly, $[z,x]-\lambda\widetilde{f_y}=-\frac{3}{\alpha-\beta}\partial_y\mathsf{w}$. Since $\mathsf{w}=\alpha\mathsf{w}_1+\beta\mathsf{w}_2+\mathsf{w}'=\frac{\beta-\alpha}{2}(\mathsf{w}_2-\mathsf{w}_1)+\frac{\beta+\alpha}{2}(\mathsf{w}_2+\mathsf{w}_1)+\mathsf{w}'$ where $\alpha,\beta\in k$ and $\mathsf{w}'\in W'$, we have $\alpha-\beta=-2\mu(\mathsf{w})$, so $\lambda=-\frac{3}{8\mu(c(\mathsf{w}))}\in k$ and $f=\overline{\mathsf{w}}+\frac{4\mu(c(\mathsf{w}))}{3}z^2\in k[x,y,z]_4$. □

5.2. **Clifford Algebras.** A 3-dimensional noetherian cubic AS-regular algebra is never a Clifford algebra in the usual sense. We need to define a new class of algebras which look like Clifford algebras.

Definition 5.2. Let $M_1,\dots,M_n\in M_n(k)$ be square matrices of degree n over k. We define a graded algebra by

$$A(M_1,\dots,M_n):=k\langle x_1,\dots,x_n,y_1,\dots,y_n\rangle/(x_ix_j^2+x_jx_ix_j+x_j^2x_i-\sum_{k=1}^{n}(M_k)_{ij}y_k;\ y_k \text{ are central})$$

with $\deg x_i=1,\deg y_i=3$.

Theorem 5.3. *Let V be a 2-dimensional vector space. If $\mathsf{w}\in\operatorname{Sym}^4 V$ is a Calabi-Yau superpotential, then $J(\mathsf{w})$ is isomorphic to $A(M_1,M_2)$ for some $M_1,M_2\in M_2(k)$.*

Proof. If $M_1=\left(\begin{smallmatrix}3&-a\\0&0\end{smallmatrix}\right),M_2=\left(\begin{smallmatrix}0&0\\-b&3\end{smallmatrix}\right)\in M_2(k)$, then $A(M_1,M_2)$ is generated by x,y,X,Y with the defining relations

$$3x^3-3X, xy^2+yxy+y^2x+aX, yx^2+xyx+x^2y+bY, 3y^2-3Y;\quad X,Y \text{ are central.}$$

It follows that

$$A(M_1,M_2)=k\langle x,y\rangle/(xy^2+yxy+y^2x+ax^3, yx^2+xyx+x^2y+by^3;\ \ x^3,y^3 \text{ are central}).$$

On the other hand, if $\mathsf{w}\in\operatorname{Sym}^4 V$ is a Calabi-Yau superpotential, then

$$J(\mathsf{w})\cong k\langle x,y\rangle/(xy^2+yxy+y^2x+ax^3, yx^2+xyx+x^2y+by^3)$$

for some $a,b\in k$ by Proposition 4.7, so it is enough to show that x^3,y^3 are central in $J(\mathsf{w})$. In $J(\mathsf{w})$,

$$\begin{aligned}
xy^3-y^3x &= -(yxy+y^2x+ax^3)y+y(xy^2+yxy+ax^3)=-ax^3y+ayx^3\\
&= ax(yx^2+xyx+by^3)-a(xyx+x^2y+by^3)x=abxy^3-aby^3x,
\end{aligned}$$

so $(1-ab)(xy^3-y^3x)$. By the classification in Proposition 4.7, $ab\neq 1$, so $xy^3=y^3x$ in $J(\mathsf{w})$. By symmetry, $yx^3=x^3y$ in $J(\mathsf{w})$, so x^3,y^3 are central in $J(\mathsf{w})$. □

Acknowledgment. The second author is indebted to Yasuhiro Ishitsuka for providing valuable information on divisors in $\mathbb{P}^1\times\mathbb{P}^1$.

References

[1] M. Artin and W. Schelter, Graded algebras of global dimension 3, *Adv. Math.* **66** (1987), 171-216.

[2] M. Artin, J. Tate and M. Van den Bergh, *Some algebras associated to automorphisms of elliptic curves*, The Grothendieck Festschrift, Vol. I, 33–85, *Progr. Math.*, **86**, Birkhauser, Boston, MA, 1990.

[3] M. Artin, J. Tate and M. Van den Bergh, Modules over regular algebras of dimension 3, *Invent. Math.* **106** (1991), 335–388.

[4] R. Bocklandt, Graded Calabi Yau algebras of dimension 3, *J. Pure Appl. Algebra* **212** (2008), 14–32.

[5] R. Bocklandt, T. Schedler and M. Wemyss, Superpotentials and higher order derivations, *J. Pure Appl. Algebra* **214** (2010), 1501–1522.

[6] K. Chan, E. Kirkman, C. Walton and J.J. Zhang, Quantum binary polyhedral groups and their actions on quantum planes, *J. Reine Angew. Math.*, in press.

[7] J. Donin and L. Makar-Limanov, Quantization of quadratic Poisson brackets on a polynomial algebra of three variables, *J. Pure Appl. Algebra* **129** (1998), 247-261.

[8] P. Jørgensen and J. J. Zhang, Gourmet's guide to Gorensteinness, *Adv. Math.* **151** (2000), 313–345.

[9] E. Kirkman and J. Kuzmanovich, Fixed subrings of Noetherian graded regular rings, *J. Algebra* **288** (2005), 463–484.

[10] E. Kirkman, J. Kuzmanovich and J. J. Zhang, Gorenstein subrings of invariants under Hopf algebra actions, *J. Algebra* **322** (2009), 3640–3669.

[11] E. Kirkman, J. Kuzmanovich, and J. J. Zhang, Shephard-Todd-Chevalley Theorem for skew polynomial rings, *Algebr. Represent. Theory* **13** (2010), 127–158.

[12] I. Mori and S. P. Smith, m-Koszul Artin-Schelter regular algebras, *J. Algebra* **446** (2016), 373–399.

[13] I. Mori and S. P. Smith, The classification of Calabi-Yau algebras with 3 generators and 3 quadratic relations, preprint.

[14] I. Mori and K. Ueyama, Ample group actions on AS-regular algebras and noncommutative graded isolated singularities, *Trans. Amer. Math. Soc.* **368** (2016), 7359–7383.

[15] M. Reyes, D. Rogalski and J. J. Zhang, Skew Calabi-Yau algebras and homological identities, *Adv. Math.* **264** (2014), 308–354.

Department of Mathematics, Graduate School of Science, Shizuoka University, 836 Ohya, Suruga-ku, Shizuoka 422-8529, Japan

E-mail address: mori.izuru@shizuoka.ac.jp

Department of Mathematics, Faculty of Education, Hirosaki University, 1 Bunkyocho, Hirosaki, Aomori 036-8560, Japan

E-mail address: k-ueyama@hirosaki-u.ac.jp